Vol. 38 No 12

BULLETIN

OF THE

TORREY BOTANICAL CLUB

DECEMBER, 1911

Dioecism in the trailing arbutus, with notes on the morphology of the seed

NEIL E. STEVENS

The heteromorphic nature of the flowers of *Epigaea repens* has been the subject of occasional remark ever since the time of André Michaux. In his journal[11] (p. 138) written while exploring in the Alleghany Mountains in 1796, occurs this entry: "Le 2 Avril Epigea repens en pleine fleur comme les jours précedents: sur plusi. individus toutes les fleurs femelles sans rudiments d'Etamines et sur d'autres individus toutes les fleurs hermaphrodites." His Flora[12] (p. 250) contains this further note: "Flores omnes in nonnullis individuis abortivi."

In 1868, Thomas Meehan[10] presented a paper before the Philadelphia Academy of Sciences in which he called attention to the fact that the pistillate flowers have "fine cleft stigmas strongly recurved, exposing a glutinous surface; while the hermaphrodite ones keep the apex of the pistils closed." From this he inferred that the plant was practically dioecious; and an examination of a number of plants during the fruiting season apparently confirmed this belief, for he found that over half the plants set no fruit.

Asa Gray[4] published in 1876 a short article on the floral structure of *Epigaea*, in which he classifies the flowers in two main groups, each with two modifications. The first group is characterized by having well developed stigmas composed of five radiate lobes which are moist and glutinous and evidently well

[The BULLETIN for November 1911 (**38**: 489-530. *pl. 27-34. f. 1*) was issued 1 D 1911.]

adapted for retaining pollen. This first group he subdivides on the basis of whether the stigma projects above the corolla or is included within the tube. In the "shorter styled form," however, the height of the stigma varies considerably, being "sometimes as low as the middle of the tube of the corolla, sometimes nearly up to the throat." This main group is further characterized by having aborted stamens. The stamens of this form show varying stages in abortion, from mere rudiments of filaments to a condition of fully developed filaments bearing anthers with occasional pollen grains. They are, however, always functionless and the form is evidently pistillate.

The second group is characterized by having small, erect, and comparatively smooth and dry stigmas which do not retain pollen readily. The ovaries in this form are apparently as well developed as those in flowers of the first group, and the styles show the same variations in length. This small stigma form has well developed stamens with anthers that abound in pollen. The stamens vary much less in length than do the pistils but there is "a tendency to having lower instead of higher anthers in connection with the short style." In conclusion Gray raises the question as to whether *Epigaea* is really dioecious or the small stigma form sometimes sets fruit; and suggests that the modifications in the length of the style point toward a heterostylous condition like that of the primroses.

Since Gray's time several papers dealing with the flowers of *Epigaea* have appeared. All of them agree in most particulars with his observations. Halsted[6] found all the kinds of flowers described by Gray, and notes that about one third of the flowers of the small stigma form have the stigmas even with the anthers. He measured the pollen grains from the longer and shorter stamens and found no difference in their size. As a difference in the size of the pollen of the different forms is common among heterostylous plants Halsted concludes that *Epigaea* shows no real evidence of heterostyly.

Wilson[15] examined a thousand plants, mostly from North Carolina. He found that the kind with perfect stigmas showed all gradations in length of style from the longest to the shortest. In only one of these large stigma plants did he find even aborted

anthers. In the kind with pollen-bearing stamens he found the same variations in length of pistil and an equal variation in length of stamen. The two organs do not show any correspondence in length, however, but all lengths of stamens are found associated with each length of pistil. Wilson concludes that the pistil in this form is functionless because of its imperfect stigma and that the species is really dioecious; and he makes the interesting observation that the flowers of the two forms differ, the corolla being from one third to one seventh smaller and generally more pink in the large stigma form than in the small stigma form. He also considers that *Epigaea* must have been trimorphic before becoming dioecious.

Miss Langdon[9] examined a large number of flowers and found all the types described above. She notes that several arrangements of anthers may occur with each length of stigma, and that while usually all the flowers of a cluster are of the same form occasional exceptions occur. For example, on one spray there was a cluster of medium-styled flowers and one of short-styled; and in one short-styled cluster was one long-styled flower. Bastin[1] in a paper on the structure of *Epigaea*, discusses the flowers but all his observations on this point are taken from Wilson's article.

The peculiar condition of the flowers of *Epigaea* was brought to the writer's attention, in connection with some work on heterostylous plants, by Gray's suggestion that *Epigaea* might be an example of a heterostylous species that either was approaching or had attained a dioecious condition. As will be noted from the papers cited above, there is considerable uncertainty as to whether *Epigaea repens* is really dioecious, that is, whether in the form having perfect stamens the pistil is incapable of producing seed; and as to whether there is any real evidence of a heterostylous condition. The writer has accordingly investigated these two points. In connection with this work, flowers from several localities in New England and New York have been examined, and a number of plants taken up and cultivated for experimental purposes. The writer wishes to acknowledge his indebtedness to friends who have sent him specimens and to Mr. F. V. Coville for generously sending information as to his methods of cultivating the plant.

EVIDENCE OF HETEROSTYLY

The claims as to heterostyly in *Epigaea* are based entirely upon the variation in the length of the pistils As Darwin[3] (p. 3) emphasized, however, in his original discussion of heterostylous plants, anatomical characters alone do not furnish conclusive evidence of heterostyly. The essential differences between the forms of a heterostylous species are physiological differences in the pistils and pollen. Unless it be proven that one form is wholly fertile only when pollinated with pollen from another form, we do not have conclusive evidence that the species is heterostyled.

In *Epigaea*, moreover, the anatomical evidence is anything but conclusive. Gray (loc. cit. 75), to be sure, divides the large stigma kind into two groups distinguished by different lengths of style, but there are all gradations from the longest style to the shortest (Wilson, loc. cit. 59). The small stigma form shows even less evidence of heterostyly, for both styles and stamens show all gradations in length and the stamens vary within much narrower limits than do the pistils (Gray, loc. cit. 75). And in place of the correlation of pistils of a given length with stamens of a different length characteristic of heterostylous flowers, we find in *Epigaea* any length of stamen associated with any length of pistil (Wilson, loc. cit. 59, and Miss Langdon, loc. cit. 11); and flowers with stigmas at the anther level are common (Halsted, loc. cit. 249). Moreover, as Halsted points out, the pollen grains from the stamens of different lengths show no difference in size; and this, while by no means conclusive, lessens the probability that the species is truly heterostylous. All these observations have been confirmed by the writer.

Experiments, however, show still more conclusively that *Epigaea* cannot be considered a heterostylous plant. The writer transplanted several wild plants with large stigma, each of which bore a dozen or more buds, and cultivated them out of doors. The plants apparently suffered no injury and all blossomed readily. While the pistils showed a wide variation in length they may be roughly classified as long-styled, mid-styled, and short-styled forms. Flowers of each of these forms were pollinated with pollen from long and mid-length stamens. No flower with stamens

corresponding in length to the shortest pistils has been found by the writer.

The pollen germinated readily, and sections made twenty-four hours after pollination showed the stigmas crowded with pollen tubes. The tubes, however, developed rather slowly, and it was only at the end of five days, 120 hours, that pollen tubes were found extending to the ovules, and in a few cases the polar nuclei appeared about to fuse with the male nucleus. It is noteworthy that the pollen from both lengths of stamens germinated with equal readiness on all lengths of pistil. Such would not be the case, of course, in a truly heterostylous plant.

DIOECISM

The evidence as to dioecism is much more satisfactory. Unquestionably the large stigma form is uniformly pistillate, for usually only mere rudiments of filaments appear and often none at all. Gray (loc. cit. 74), however, found in some anthers in a flower of this form a few "perhaps well formed pollen grains," but the anthers never opened. In the one thousand plants examined by him, Wilson (loc. cit. 59) found only one large stigma plant that showed even aborted anthers. The writer examined over two hundred plants of the large stigma form, and only three bore flowers with any rudiments of anthers. Microtome sections of these flowers were prepared, and while the four locules could in some cases be distinguished they contained merely masses of broken down cells.

The form with well developed stamens, on the other hand, has never been observed to set fruit; and as Gray (loc. cit. 76) points out, it is easy to ascertain in any case which kind of flower has matured fruit, because the style and stigma persist until the fruit is fully mature. Indeed, in the flowers in which the ovary has not developed, the style and stigma often persist for over a year, somewhat shrunken, to be sure, but still sufficiently well preserved to make it certain to which form the flower belongs. This unusual persistence of the style and stigma seems to be due to the fact that the carpellary bundles contain strongly lignified tracheids.

The pistil of this form, however, is, as Gray (loc. cit. 75) observed, apparently normal except for its smaller and rather

dry stigma. The writer experimented with the artificial pollination of this small stigma form both with plants cultivated in pots and with flowers kept in water. The pollen adheres to these stigmas to a considerable extent, though this is probably due to the fact that the wall of the pollen grain is somewhat mucilaginous, for the pollen adheres also to the style below the stigma and even to smooth objects, such as a glass rod. In no case, however, was a pollen grain observed to develop a tube when placed on a stigma of this kind. The pollen was retained without degeneration for some time, frequently for over a week, but it never established any organic connection with the pistil and could easily be brushed off. As these cultivated plants appeared perfectly healthy and afterward developed apparently normal new shoots, the failure of the pollen to germinate cannot be attributed to any unusual weakened condition of the pistils. Moreover, the same condition was found to occur in long-styled plants of the small stigma form growing under natural conditions. Although the stigma in these plants projected well above the stamens it was covered with pollen, probably as the result of the visits of insects. Not a single grain, however, had emitted a tube.

This evident sterility of the pistils in the staminate form seemed still more remarkable when microtome sections of the ovules were examined. At the time of fertilization the embryo sac of the large stigma form shows uniformly typical egg and synergids, well defined antipodal cells, and the two polar nuclei lying close together near the middle of the sac. In all probability it remains in this condition over winter (Coulter and Chamberlain,[2] p. 53). The ovules of the sterile small stigma form present an identical appearance both in size and in the condition of the embryo sac. It is of course possible that the nuclei of this functionless embryo sac have not undergone a reduction division. There appears to be no evidence, however, of a tendency toward apogamy, and these unfertilized ovules gradually degenerate.

We have then, in the small stigma form of *Epigaea*, flowers that are apparently perfect but functionally male. The morphological differences between the functional and non-functional pistils in this species are, as pointed out above, very slight and are confined to the stigmas. Their physiological differences are so

great, however, that pollen which germinates readily on the stigmas of one form is absolutely inert on those of the other, and *Epigaea repens* is functionally a dioecious plant.

A few other cases of plants that are apparently polygamodioecious but actually dioecious have been observed. Darwin (loc. cit. 288) noted that the "hermaphrodite" flowers of *Euonymus europaeus* are practically male, so this species is really dioecious. Mottier[13] (p. 377) and von Kirchner[8] (p. 116) have pointed out that the apparently polygamodioecious maples are functionally dioecious, because the anthers of the "perfect" flowers, although they contain some apparently normal pollen grains, never open.

Celastrus scandens is an apparently polygamodioecious plant, having pistillate and "perfect" flowers. Like the related *Euonymus europaeus*, however, it is dioecious in function, for the pistils of the "perfect" flowers are smaller than those of the pistillate and appear upon investigation to be non-functional. *Celastrus* is abundant in the vicinity of New Haven, Connecticut, and during the spring of 1911 the writer examined a large number of plants in this region. Particular care was taken to observe the plants bearing "perfect" flowers. Not only was there no indication of any development of the ovary but the flowers shriveled and dropped from the plant almost as soon as the pollen was shed. Microtome sections showed that the ovules in the pistillate flowers are more than ten times as large as those of the "perfect" flowers. Many of these non-functioning ovules contain, however, an apparently typical embryo sac.

The condition found in the maples is exactly the reverse of that found in *Epigaea* and *Celastrus*. In the latter species there are two kinds of flowers, one evidently pistillate, the other developing both ovules and pollen grains. In some species of maples, on the other hand, one kind of flower is evidently staminate and the other apparently perfect. In *Epigaea* and *Celastrus*, however, the ovules of the "perfect" flowers are never fertilized, and in the maples the anthers of the "perfect" flowers do not open. Notwithstanding their morphological differences, then, both types are functionally dioecious.

As Wilson (loc. cit. 59) states, there is in *Epigaea repens* an apparently constant difference in size between the large-stigma

and small-stigma flowers. If flowers of the two forms growing in the same locality are compared, the pistillate flowers seem to be uniformly smaller than the apparently perfect flowers. Darwin (loc. cit. 307) considers that this is generally the case in gynodioecious plants, and names a dozen such species in which the pistillate flowers are smaller than the staminate flowers. He notes also the fact that no such difference in the size of the corolla has been found in the two forms of androdioecious species.

The writer has been unable to demonstrate, however, any correlation of color of the corolla with the two forms of stigma. Wilson (loc. cit. 59) states that the corollas of the large stigma flowers are pink, while those of the small stigma flowers are white. This is apparently true in some localities; but in one lot of flowers sent from a locality in Maine nearly all the flowers of both forms were pink. Moreover, a small lot of flowers from New Hampshire showed the large stigma flowers small and white, while the small stigma flowers were larger and very deep pink.

MORPHOLOGY OF THE SEED

In the sections made for the purpose of determining the rate of growth of the pollen tubes several points were noted which seemed worthy of record, and the development of the seed has



FIG. 1. Longitudinal section through ovule of small stigma form of *Epigaea repens* showing "tapetum" surrounding the lower portion of the embryo sac, elongated egg and synergid, saclike antipodal cells and polar nuclei. This ovule though non-functional is identical in appearance with the functional ones. ×400.

accordingly been traced. The mature embryo sac (FIG. 1) presents no unusual features except that it is somewhat curved and is prolonged, toward its micropylar end, into a narrow funnel-shaped region which contains the greater part of the egg and synergids. The nucellus either does not persist or else it coalesces so completely with the integument that they appear continuous.

The single massive integument, which makes up the bulk of the ovule, is differentiated into three rather definite regions: the epidermis, a middle storage region, and an inner tapetal region surrounding the embryo sac. The cells of the epidermis are large, regular, and rather thick-walled. The storage region shows little differentiation except that the cells in the funicular region are considerably elongated. The "tapetum" is composed of a single layer of cells except at the antipodal end, where it is usually two or more cells in thickness. Its cells have the large nuclei and dense granular cytoplasm characteristic of nutritive cells. It is noteworthy that the tapetum does not surround the narrow micropylar portion of the embryo sac, a fact which suggests that the micropylar portion may be a secondary development due to an encroachment of the growing egg and synergids upon the tissues of the integument.

As noted above, at the time of maturity of the embryo sac the polar nuclei lie close together in the middle of the sac. They remain entirely distinct, however, until fertilization occurs and the primary endosperm nucleus is formed by the simultaneous fusion of the polar nuclei with the male nucleus. The division of the primary endosperm nucleus is accompanied by the formation of a transverse wall dividing the embryo sac into two approximately equal chambers (FIG. 2). Each of these chambers is further divided by a cross wall resulting in a "four-chambered" embryo sac (FIG. 3). No division of the fertilized egg has occurred at this stage.

In the further development of the endosperm, cell divisions occur rapidly in all four of these "chambers." The cell divisions take place in various planes, and by the time the proembryo contains five or six cells it is impossible to distinguish the original cross walls which separated the four chambers of the embryo sac. There is thus no period of free nuclear division in the development of the endosperm in *Epigaea*.

Instances in which the first division of the primary endosperm nucleus is followed by the formation of a transverse wall across the embryo sac, are fairly common especially among the Sympetalae (Coulter and Chamberlain, loc. cit. 176); and the formation of a two-chambered embryo sac has been reported in the three subfamilies most closely related to the Ericoideae, that is,



FIGS. 2 and 3. "Chambered" embryo sac of *Epigaea repens*. ×600.

FIG. 2. Two-celled stage.

FIG. 3. Four-celled stage. The fertilized egg is partly covered by the remains of the pollen tube, and the antipodal cells have largely degenerated.

in the Pyroloideae, Monotropoideae, and Vaccinioideae (Hofmeister,[7] p. 141). The later development of the endosperm differs somewhat in the different groups. In the Vaccinioideae the endosperm is reported to develop only in the antipodal chamber while in the Pyroloideae and Monotropoideae endosperm is formed in both chambers. The only case in which, as in *Epigaea*, the embryo sac becomes divided into four superposed chambers all

of which take part in the development of the endosperm, is that of *Datura laevis* reported by Guignard[5] (p. 166).

A " four-chambered " embryo sac, formed, however, somewhat differently from those of *Datura* and *Epigaea*, is found in *Physostegia virginiana*. In this species Sharp[14] (p. 220) has recently observed that the division of the primary endosperm nucleus is accompanied by the formation of a longitudinal wall running through the middle of the embryo sac. The nuclei of the two resulting parts of the sac then divide and transverse walls are



FIG. 4. Longitudinal section of developing ovule of *Epigaea repens*, showing haustoria which function for the nourishment of the endosperm. ×190.

formed. Further transverse divisions then give rise to a large-celled thin-walled tissue which fills the " endosperm lobe."

By the time the embryo of *Epigaea* reaches the quadrant stage the endosperm has a very considerable bulk and shows in longitudinal section fifty or more cells. At about this stage a knobshaped projection appears at either end of the endosperm. These outgrowths extend into the tissues of the integument and apparently function as haustoria to nourish the developing endosperm.

At first both haustoria consist of but a single large cell; and this is usually the condition of the antipodal haustorium at

maturity. In the formation of this cell a portion of one of the peripheral cells of the endosperm grows out into the tissues of the integument, increases greatly in size, and is later cut off by a cross wall. By the continual growth and division of the cell thus cut off, the micropylar haustorium develops rapidly until finally it consists of a group of eight or ten cells having dense granular contents and large nuclei. Frequently some of the cells of this haustorium contain two nuclei. The condition shown in FIG. 4 represents about the maximum development of the haustoria, and in the nearly mature seed they appear crushed and distorted among the tissues of the integument.

Haustoria of various kinds that function for the nourishment of the endosperm have been reported in numerous instances. Their occurrence and origin, together with the whole subject of the nourishment of the developing embryo sac, have been discussed by Coulter and Chamberlain (loc. cit. 104–113). They mention haustoria of various forms, developed from the micropylar and chalazal ends and even from the sides of the sac. Sometimes the haustoria develop from protrusions of the sac itself; but in other forms the synergids, the antipodal cells, the suspensor, and one of the row of megaspores have been reported to take part in the formation of haustoria. The structures found in *Epigaea* are noteworthy as rather simple haustoria which originate directly from the endosperm at a comparatively late stage in its development.

BOTANICAL LABORATORY,
YALE UNIVERSITY.

LITERATURE CITED

1. **Bastin, E. S.** Structure of *Epigaea repens*. Am. Jour. Pharm. **67**: 231–236. 1895.
2. **Coulter, J. M., & Chamberlain, C. J.** Morphology of angiosperms. New York, 1903.
3. **Darwin, C.** The different forms of flowers on plants of the same species. New York, edition of 1897.
4. **Gray, A.** Heteromorphism in *Epigaea*. Am. Jour. Sci. III. **12**: 74–76. 1876. (Same article) Am. Nat. **10**: 490–492.
5. **Guignard, L.** La double fécondation chez les Solanées. Jour. de Bot. **16**: 145–167. 1902.

6. **Halsted, B. D.** Notes upon *Epigaea repens*. Bull. Torrey Club **18**: 249, 250. 1891.
7. **Hofmeister, W.** Neuere Beobachtungen über Embryobildung der Phanerogamen. Jahrb. Wiss. Bot. **1**: 82–188. 1858.
8. **Kirchner, O. von.** Blumen und Insekten. Leipzig, 1911.
9. **Langdon, F. E.** A study of *Epigaea repens*. Asa Gray Bull. No. 4, First Quarter, 1894.
10. **Meehan, T.** *Mitchella repens*, a dioecious plant. Proc. Acad. Nat. Sci. Phila. **1868**: 153–156, 183–184.
11. **Michaux, A.** Extracts from the journal of André Michaux written during his travels in the U. S. and Canada, 1785–1796. Proc. Am. Philos. Soc. **26**: 1–145. 1889.
12. **Michaux, A.** Flora Bor.-Am. Paris, edition of 1820.
13. **Mottier, D. M.** Development of the embryo sac in *Acer rubrum*. Bot. Gaz. **18**: 375–377. 1893.
14. **Sharp, L. W.** The embryo sac of *Physostegia*. Bot. Gaz. **52**: 218–224. 1911.
15. **Wilson, W. P.** Observations on *Epigaea repens*. Contr. Bot. Lab. Univ. Penn. **1**: 56–63. 1893.

On the identity of Cyathea multiflora, type of the genus Hemitelia R. Br.*

WILLIAM R. MAXON

(WITH PLATE 35)

The genus *Hemitelia* was established by Robert Brown in 1810 for a group of tree ferns, of which the following previously described species were known to him and were stated as representatives of his new genus: (1) *Cyathea multiflora* Sm., (2) *C. horrida* Sm., and (3) *C. capensis* Sm. Fortunately these are now regarded, as they have been generally in the past, as congeneric. If, then, the genus *Hemitelia* is to be recognized as valid, there can be no objection to a selection of the first-named species, *Cyathea multiflora*, as its type. This course has been followed in recent years by both L. M. Underwood† and C. Christensen,‡ and has been tacitly accepted, although quite a different typification of the genus was long ago proposed by G. Gardner.§

Concerning the identity of *Cyathea multiflora*, however, there has been little but confusion, almost from the time of its publication. Following a brief description (quoted from Willdenow) Hooker commented upon this species in the Species Filicum‖ as follows:

> "Hab. Jamaica; ex Herb. Banks. (*Smith*).—With this I am unacquainted, and Mr. Brown is the authority for its being referred to *Hemitelia*. Sir Jas. Smith, with whom the species originated, has merely said of it (under *Cyathea*) 'Caudice ——, fronde bipinnata pinnatifida, laciniis obtusis serratis, rachi alata, floribus sparsis, calyce lacero.' Willdenow, who seems to have been acquainted with the species, and whose character I have given above, further remarks upon it, 'Rachis margined on each side with a narrow decurrent line. Partial pinnæ 2 feet long. Pinnules 4 inches long, an inch broad at

* Published by permission of the Secretary of the Smithsonian Institution.

† Mem. Torrey Club **6**: 268. 1899.

‡ Index Fil. XVII. 1906.

§ London Journ. Bot. **1**: 438–442. 1842.

‖ Sp. Fil. **5**: 32. 1846.

the base, lanceolate, acuminate, pinnatifid. Segments 5 lines long, oblong, rather acute, obtusely serrated.' Unfortunately neither Smith nor Willdenow alludes to the venation, nor, except the brief notice of the former, to the involucres. Mr. Smith, who, as well as Mr. Gardner, has examined the original specimens, says that, as far as can be judged from the imperfect specimens, it differs from the preceding (*H.? Parkeri*) only in wanting the coarse hairs on the rachis. Mr. Gardner considers it allied to *Alsophila capensis*."

It may here be noted that Willdenow in all probability had not seen the original specimens of the species. His description,* which very greatly modifies and extends Smith's original diagnosis, is thus one of the earliest misapplications of the species-name and is apparently the principal false basis from which so great an amount of misunderstanding has since arisen.

Smith's comment upon the source of his specimens, " Habitat in Jamaica; ex herb. Banks," led Dr. Underwood to look up the actual type of the species in the collection of Sir J. E. Smith at the rooms of the Linnean Society of London, a few years ago. He examined also the original specimens in the Banksian Herbarium (now preserved in the British Museum, Natural History), from which the three pinnæ of Sir J. E. Smith's herbarium had been derived. His conclusion was that the specimens of the two collections, which are identical, represent a species otherwise unknown. Concerning the original source of the material he has written (in MS.), " Jamaica, *R. Shakespeare*. Known only from plants collected in Jamaica in the last of the 18th century."

More recently the writer has received through the kindness of A. B. Rendle, F.L.S., Keeper of the Herbarium of the British Museum, a photograph of Banks's original specimens in the British Museum, here shown in PLATE 35, and with this two pinnules of the plant itself. A transcript of the data attached to the specimen, as shown by the photograph, is as follows:

" Shakespeare, Robert, (fl. 1780–82). Collected in Jamaica (Hb. Banks).—10. 7. 1903. Prof. L. M. Underwood called to see this type (?); hence the difficulties.— ? Can this specimen be the *type* of Hemitelia multiflora R. Br. Fl. Nov. Holl. p. 158.

* Sp. Pl. 5: 496. 1810.

= *Cyathea multiflora* Sir J. E. Smith in Act. Taur. *5*. p. 416.
= *Amphicosmia multiflora* G. Gardner in London Journ. of Bot. *1*. 1842, p. 441. Cf. this last reference, pp. 438–442. Our plant agrees with Gardner's description, p. 439.—
But Sir J. E. Smith says ' Jamaica, ex herb. Banks ' (fide Hook. Sp. Fil. *1*. 32), while *our specimen* bears '*America merid. R. Shakespeare.*' And no other specimen appears to have come from Jamaica. Shakespeare did collect in Jamaica about 1780. If this is the type, the locality ' Jamaica ' may be at fault."

From a careful comparison of the two pinnules sent, and of the photograph, with the large series of specimens in the U. S. National Herbarium, the following statements may safely be made:

1. The original specimens in the Banks Collection, British Museum, from which the three type pinnæ of *Cyathea multiflora* J. E. Smith were taken, are from a rather small frond of a species described later upon Central American material as *Hemitelia nigricans* Presl.

2. Smith's statement of the type locality of *C. multiflora* as " Jamaica " is without much doubt erroneous, since: (*a*) The original Banks specimens are marked as from " America merid."; (*b*) in all the botanical exploration carried on subsequently in Jamaica the species has not been there collected; (*c*) *Hemitelia nigricans* is a widespread though not very common species of the Atlantic coast of the mainland, extending from Guatemala, through Nicaragua to Costa Rica and Panama (and in all probability farther, along the northern coast of South America), and may reasonably be supposed to include specimens gathered by chance at almost any point along the Atlantic coast mentioned. Even if there were no confusion as to the locality for the original specimens of *C. multiflora*, their identity with *H. nigricans* can not be questioned, as the following notes are intended to make clear.

Hemitelia nigricans was described in 1849* by Presl from specimens said to have been collected by Friedrichsthal " upon the banks of the river San Juan, in Guatemala." Fournier, however,

* Presl, Epim. Bot. 31. 1849.

has stated* that the locality is in Nicaragua. There is at hand a pinnule from Presl's type at Prag, and also a pinnule of Levy's 468 from Nicaragua, mentioned by Fournier. A close comparison of these with the pinnules of *C. multiflora* from the Banks herbarium shows all three collections to be of the same species and, curiously enough, to give no indication of the variation in size and venation which exists in the more ample material of later collections. In all three the veins are almost without exception simple, and the pinnules only from 3 to 4 cm. long. Recent specimens show pinnules up to 7 cm. long and many of the veins to be once forked at or beyond the middle. From minute characters, nevertheless, it is apparent that all are referable to a single species, which naturally must bear the name *Hemitelia multiflora*. The species has since been described from Costa Rican specimens as *Alsophila decussata* Christ. It is doubtful also if *Hemitelia Hartii* Baker,† from Panama, can be maintained as a distinct species. *Hemitelia obscura* Mett.‡ is a related species described from Colombia, very similar in form, but differing conspicuously in having the lower surface of the costae strongly hispid.

From the whole series at hand, but especially from specimens collected recently by the writer in Panama, the following description has been drawn:

HEMITELIA MULTIFLORA (J. E. Smith) R. Br.; Spreng. Syst. Veg. **4**: 126. 1827

Cyathea multiflora J. E. Smith, Mem. Acad. Tur. **5**: 416. 1793.
Amphicosmia multiflora Gardner, London Journ. Bot. **1**: 441. 1842.
Hemitelia nigricans Presl, Epim. Bot. 31. 1849.
Alsophila decussata Christ in Pittier, Prim. Fl. Costar. **3**: 41. 1901.

Caudex erect, 1 to 5 meters high, very slender (about 3.5 cm. in diameter), usually sheathed by the closely ascending or semiadnate fibrous bases of old fronds, or these slowly deciduous, the scars spaced, oblong-elliptic, 4 to 5 cm. long, less than 1.5 cm.

* Bull. Soc. Bot. France **19**: 261. 1872.

† Journ. Bot. **24**: 243. 1886.

‡ *Hemitelia obscura* Mett. Ann. Sci. Nat. V. **2**: 264. 1864. Subsequently figured by Karsten, Fl. Colomb. **2**: *pl. 197. f. 2.* 1869.

broad; fronds spreading, 1.3 to 2.1 meters long; stipe up to 45 cm. long, curved, light brown, sulcate ventrally, noticeably brownish-pubescent, freely armed with straight or slightly curved spreading spines 3 to 5 mm. long, and clothed toward the base with deciduous ovate-lanceolate attenuate glossy bright-brown scales 1.5 to 2 cm. long with narrowly erose paler margins, similar scales rather densely clothing the crown and upper part of the caudex; lamina 1 to 1.7 meters long, 60 to 100 cm. broad at the middle, ovate, deeply tripinnatifid, abruptly acuminate, the primary rachis scantily aculeate near the base, upward sparsely muricate or smooth, stramineous-brownish, antrorsely brownish-strigose upon the upper side, below closely beset with stiffish mainly retrorse subpersistent yellowish hairs; pinnæ articulate, 14 to 17 pairs, approximate, the basal ones horizontal or usually deflexed, opposite and relatively short (25 to 32 cm. long, 8 to 11 cm. broad), elliptic-oblong, petiolate (1.5 to 3 cm.); middle pinnae larger, alternate, widely spreading, exactly linear-oblong, not narrowed at the base, 30 to 50 cm. long, 7 to 14 cm. broad, sessile or subsessile, abruptly acuminate-caudate, the secondary rachis like the primary, in the outer part delicately foliaceo-marginate; pinnules 20 to 24 pairs, articulate, deciduous, mostly spreading at right angles, slightly separated or half their width apart, all but the lowermost strictly sessile, linear-oblong, 3 to 7 cm. long, 9 to 18 mm. broad, acutish or rarely acuminate, regularly pinnatifid two thirds to four fifths the distance toward the slender elevated costa; costae yellowish-strigose above, below bearing a few minute deciduous yellowish brown subbullate attenuate scales, glabrous or minutely setulose; segments 9 to 12 pairs, slightly oblique, straight or slightly falcate, oblong, close, the margins plane or lightly revolute, crenate-dentate, strongly so at the very obtuse or sometimes muticous apex, the sides nearly parallel; costules bearing 2 to 4 stiff spinelike hairs above, below glabrous or obscurely and minutely setulose; veins wide-spreading, 4 to 8 pairs, simple or once forked at or beyond the middle, dark, more or less concealed, extending to the margin; sori 1 to 4 pairs, mostly medial, the lower ones supramedial, borne at or just below the fork; indusium proximal, brown, membranous, cucullate-dimidiate, finally somewhat reflexed and cleft into 2 or 3 shallow lobes, often nearly disappearing with age; receptacle globose, the numerous flaccid paraphyses equaling the sporangia; leaf tissue papyraceo-herbaceous, translucent, very dark green above, much paler below, glabrous.

TYPE LOCALITY: "America meridionalis," *R. Shakespeare.*

DISTRIBUTION: Guatemala, Nicaragua, Costa Rica, and

Panama, ascending to 1,100 meters altitude; also reported from Peru.

Besides the fragment of the Shakespeare type above mentioned the following specimens are in the U. S. National Herbarium:

NICARAGUA: Banks of the Rio San Juan, *Friedrichsthal.* Chontales, in forest, *Levy* 465.

GUATEMALA: Forest near Livingston, at sea-level, June, 1906, *H. von Türckheim* II. 1221. Same locality, February 18, 1905, *C. C. Deam* 483 (determined as *Cyathea arborea*). Same locality, January 18, 1905, *Kellerman* 5080.

COSTA RICA: Forêt à Général, altitude 600 meters, *Pittier* 3344. Fougeraies du Général, vallée du Diquís, altitude 700 meters, *Pittier* 12027 (type collection of *Alsophila decussata* Christ). Cienegas de Cañas Gordas, *Pittier* 10978. Cañas Gordas, altitude 1,100 meters, *Pittier* 10980.

PANAMA: Laguna de Chiriqui and its neighborhood, *J. Hart* 12, 29. Humid forest near Porto Bello, province of Colon, at sea-level, April 7, 1911, *Maxon* 5776.

Finally, it may be pointed out that the small size of the original specimens of *H. multiflora*, and perhaps also of Presl's type of *H. nigricans*, is further due to their having come from the upper part of a small frond. The type fragments of both are exactly matched by pinnules from the apical pinnae of the writer's specimens from Panama (the only apical portions at hand), in which also two other perfectly diagnostic features are evident, namely, the conspicuously winged secondary rachises, and the peculiar long-acuminate apices of the pinnae. Both may be readily observed in the illustration herewith published.

WASHINGTON, D.C.

Explanation of plate 35

From a photograph of the original specimens in the Banksian Herbarium, British Museum (Natural History), from which the type of *Cyathea multiflora* was derived. Collected in "America meridionalis" by R. Shakespeare. About one third natural size.

INDEX TO AMERICAN BOTANICAL LITERATURE

(1905-1911)

The aim of this Index is to include all current botanical literature written by Americans, published in America, or based upon American material; the word America being used in the broadest sense.

Reviews, and papers that relate exclusively to forestry, agriculture, horticulture, manufactured products of vegetable origin, or laboratory methods are not included, and no attempt is made to index the literature of bacteriology. An occasional exception is made in favor of some paper appearing in an American periodical which is devoted wholly to botany. Reprints are not mentioned unless they differ from the original in some important particular. If users of the Index will call the attention of the editor to errors or omissions, their kindness will be appreciated.

This Index is reprinted monthly on cards, and furnished in this form to subscribers at the rate of one cent for each card. Selections of cards are not permitted; each subscriber must take all cards published during the term of his subscription. Correspondence relating to the card issue should be addressed to the Treasurer of the Torrey Botanical Club.

Andrews, A. L. Notes on North American *Sphagnum*—I. Bryologist **14**: 72–75. Jl 1911.

Appleman, C. O. Physiological behavior of enzymes and carbohydrate transformations in after-ripening of the potato tuber. Bot. Gaz. **52**: 306–315. 17 O 1911.

Arechavaleta, J. Flora uruguaya **4**: 129–224. *f. 1–14.* 1911. An. Mus. Nac. Montevideo vol. 7.

Includes *Ipomoea biglandulosa, I. pinifolioides, I. tacuaremboense* spp. nov.

Atwood, A. C. Description of the comprehensive catalogue of botanical literature in the libraries of Washington. U. S. Dept. Agr. Plant Ind. Circ. 87: 1–7. 9 S 1911.

Bartlett, H. H. Systematic studies on *Oenothera*—I. *Oenothera Tracyi* sp. nov. Rhodora **13**: 209–211. *pl. 93.* 9 O 1911.

Bancroft, K. A note on the canker of *Hevea brasiliensis.* Agric. Bull. Straits and Fed. Malay States **10**: 203–208. Jl 1911.

Blanchard, W. H. *Lycopodium complanatum* near Hartland, Vermont. Rhodora **13**: 211, 212. 9 O 1911.

Blanchard, W. H. *Rubus* of eastern North America. Bull. Torrey Club **38**: 425–439. 6 O 1911.

Berry, E. W. Contributions to the Mesozoic flora of the Atlantic

coastal plain—VII. Bull. Torrey Club **38**: 399–424. *pl. 18, 19.* 6 O 1911.

Includes descriptions of 10 new species and 2 new genera.

Bödeker, F. *Mamillaria Ottonis* Pfeiff., *Mam. Golziana* Ferd. Haage jun. und *Mam. Bussleri* Mundt. Monats. Kakteenk. **21**: 140–142. 15 S 1911.

Brand, C. J. Grimm alfalfa and its utilization in the Northwest. U. S. Dept. Agr. Plant Ind. Bull. 209: 1–66. *pl. 1, 2.* 27 My 1911.

Bresadola, G. Fungi borneenses. Ann. Myc. **9**: 549–553. 1 O 1911.

Appendix contains 8 fungi from Trinidad.

Brown, W. H. The development of the ascocarp of *Lachnea scutellata*. Bot. Gaz. **52**: 275–305. *pl. 9+f. 1–51.* 17 O 1911.

Cardiff, J. D. An aberrant walnut. Trans. Kansas Acad. Sci. **23, 24**: 138. Ja 1911.

Cardot, J. Coup d'oeil sur la flore bryologique du Mexique. Rev. Bryol. **38**: 79–84; 97–108. 1911.

Cary, M. A biological survey of Colorado. N. Am. Fauna **33**: 1–256. *pl. 1–12+f. 1–39.* 17 Au 1911.

Contains lists of plants found in the various life zones.

Christensen, C. Revision of the American species of *Dryopteris* of the group of *D. opposita*. Danske Vid. Selsk. Skr. VII. Nat. **4**: 249–336. 1907.

Clute, W. N. The distribution of *Asplenium alternans*. Fern Bull. **19**: 38–42. [O] 1911.

Clute, W. N. *Hemigramma latifolia*. Fern Bull. **19**: 46–48. [O] 1911. [Illust.]

[Clute, W. N.] Pteridographia. Fern Bull. **19**: 53–56. [O] 1911.

Includes notes on (*a*) forked liquorice fern, (*b*) sex in the ostrich fern, (*c*) cambium in ferns, (*d*) fragrant bracken, and (*e*) abnormal lycopodiums.

[Clute, W. N.] Rare forms of ferns—XVIII. Fern Bull. **19**: 50–52. [O] 1911. [Illust.]

Collins, G. N. Increased yields of corn from hybrid seed. Yearbook U. S. Dept. Agr. **1910**: 319–328. 1911.

Cook, O. F. Hindi cotton in Egypt. U. S. Dept. Agr. Plant. Ind. Bull. 210: 1–58. *pl. 1–6.* 11 My 1911.

Cook, O. F., & Meade, R. M. Arrangement of parts in the cotton plant. U. S. Dept. Agr. Plant Ind. Bull. 222: 1–26. *f. 1–9.* 3 O 1911.

Corbett, L. C. The school garden. U. S. Dept. Agr. Farm. Bull. 218: 1–40. *f.* 1–33. 1905.

Spanish translation in Republica de Colombia Revista del Ministerio **5**: 13–44 1911.

Dallimore, W. Notes on trees suitable for experimental forestry—II. American broad-leaved trees. Kew Bull. Mis. Inf. **1911**: 303–310. O 1911.

Damianovich, H. La doctrina de la generación espontánea: su evolución y estado actual. Anal. Soc. Ci. Argent. **71**: 153–192. *f.* 1–10. Ap 1911; **71**: 193–230. *f.* 11–22. My 1911.

Davis, J. J. Fourth supplementary list of parasitic fungi of Wisconsin. Trans. Wisconsin Acad. **16**: 739–772. [1910.]

Davy, J. B. Observations on the inheritance of characters in *Zea Mays* Linn. Science II. **34**: 576. 27 O 1911.

Diels, L. *Menispermaceae.* Pflanzenreich 4^{94}: 1–345. *f.* 1–93. 6 D 1910.

Dominguez, J. A. Datos para la materia médica Argentina **2**: i–ix +1–141. Buenos Aires, 1910.

Constitutes no. 25 of Trab. Inst. Farm Fac. Cien. Med. Buenos Aires.

Dox, A. W., & Golden, R. Phytase in lower fungi. Jour. Biol. Chem. **10**: 183–186. O 1911.

Eichlam, F. Mitteilungen aus Zentral-Amerika—IX. Monats. Kakteenk. **21**: 113–116. 15 Au 1911;—X. Monats. Kakteenk. **21**: 129–131. 15 S 1911.

Evans, A. W. Notes on North American *Hepaticae*—II. Bryologist **14**: 84–88. S 1911.

Fairchild, D. Seeds and plants imported during the period from January 1 to March 31, 1910. U. S. Dept. Agr. Plant Ind. Bull. 207: 1–100. 4 My 1911.

Fairchild, D. Seeds and plants imported during the period from April 1 to June 30, 1910. U. S. Dept. Agr. Plant Ind. Bull. 208: 1–88. 29 Ap 1911.

Fawcett, H. S., & Burger, O. F. A variety of *Cladosporium herbarum* on *Citrus aurantium* in Florida. Phytopathology **1**: 164–166. O 1911.

Fedde, F. *Papaveraceae-Hypecoideae* et *Papaveraceae-Papaveroideae.* Pflanzenreich 4^{104}: 1–430. *f.* 1–43. 21 D 1909.

Fernald, M. L. Harshberger's phytogeographic survey of North America. Rhodora **13**: 213–224. 9 O 1911.

Contains much original matter on the distribution of plants.

Ferriss, J. H. The finding of *Asplenium alternans*. Fern Bull. **19**: 36–38. [O] 1911.

Fink, B. The nature and classification of lichens—I. Views and arguments of botanists concerning classification. Mycologia **3**: 231–269. S 1911.

Fletcher, E. F. *Dracocephalum thymiflorum* a casual plant at Westford, Massachusetts. Rhodora **13**: 212. 9 O 1911.

Foxworthy, F. W. Bedaru and Billian: two important Borneo timber trees. Philip. Jour. Sci. **6**: (Bot.) 179, 180. Jl 1911.
Urandra corniculata and *Eusideroxylon Zwageri*.

Foxworthy, F. W. Philippine *Dipterocarpaceae*. Philip. Jour. Sci. **6**: (Bot.) 231–287. *pl. 34–44*. S 1911.
Four new species described.

Freeman, E. M., & Johnson, E. C. The rusts of grains in the United States. U. S. Dept. Agr. Plant Ind. Bull. 216: 1–87. *pl. 1+f. 1, 2*. 15 Au 1911.

Fries, R. E. Entwurf einer Monographie der Gattungen *Wissadula* und *Pseudabutilon*. Kgl. Sv. Vet.-Ak. Handl. 43^4: 1–114. *pl. 1–10*. 1908.
The new species of *Wissadula* are also published in Repert. Sp. Nov. **9**: 486–492. 15 Au 1911; **9**: 504–513. 5 S 1911.

Fries, R. E. Studien über die amerikanische Columniferenflora. Kgl. Sv. Vet.-Ak. Handl. 42^{12}: 1–67. *pl. 1–7*. 1908.
Includes 21 new species.

Gates, F. C. A bog in central Illinois. Torreya **11**: 205–211. *f. 1–3*. 18 O 1911.

Gates, R. R. Mutation in *Oenothera*. Am. Nat. **45**: 577–606. O 1911.

Gilbert, E. M. Studies on the *Tremellineae* of Wisconsin. Trans. Wisconsin Acad. **16**: 1137–1170. *pl. 82–84*. O 1910.

Greenwood, H. E. Some stages in the development of *Pellia epiphylla*. Bryologist **14**: 59–70. *pl. 7–9*. Jl 1911; **14**: 77–83. *pl. 10–12*. S 1911.

Grout, A. J. Notes on Vermont bryophytes—VI. Bryologist **14**: 52–54. 19 My 1911.

Guillaumin, A. L'étude des germinations appliquée à la classification des genres et à la phylogénie des groupes. Rev. Gén. Bot. **22**: 449–468. *pl. 12, 13*. 15 D 1910.

Hall, H. M. Additional southern Californian *Compositae*. Muhlenbergia **7**: 83–84. 30 S 1911.

Hassler, E. Contribuciones à la flora del Chaco Argentino-Paraguayo. Primera parte: Florula pilcomayensis. i–iii+1–154. Buenos Aires, 1909.
Constitutes no. 21 Trab. Mus. Farm. Fac. Cien Méd. Buenos Aires.

Heald, D., & Wolf, F. A. List of parasitic bacteria and fungi occurring in Texas. Trans. Texas Acad. Sci. **11**: 10–44. 1911.

Hedgcock, G. G. The cross-inoculation of fruit trees and shrubs with crown-gall. U. S. Dept. Agr. Plant Ind. Bull. 131: 21–23. 16 Mr 1908.

Hedgcock, G. G. Field studies of the crown-gall and hairy-root of the apple tree. U. S. Dept. Agr. Plant Ind. Bull. 186: 1–108. *pl. 1–10.* 17 N 1910.

Hedgcock, G. G. Field studies of the crown-gall of the grape. U. S. Dept. Agr. Plant Ind. Bull. 183: 1–40. *pl. 1–4.* 23 Jl 1910.

Hedgcock, G. G. Some stem tumors or knots on apple and quince trees. U. S. Dept. Agr. Plant Ind. Circ. 3: 1–16. *f. 1–10.* 11 My 1908.

Henslow, G. The origin and history of our garden vegetables and their dietetic values—II. Jour. Royal Hort. Soc. **36**: 345–357. *f. 120–125.* N 1910.
Roots and tubers.

Heese, E. *Echinocactus Gürkeanus* Heese, nov. spec. Monats. Kakteenk. **21**: 132, 133. 15 S 1911. [Illust.]

Heller, A. A. The North American lupines—IV. Muhlenbergia **7**: 13–15. *f. 5.* 14 Mr 1911.
Lupinus apodotropis sp. nov.

Hemenway, A. F. Studies on the phloem of the dicotyledons—I. Phloem of the *Juglandaceae.* Bot. Gaz. **51**: 131–135. *pl. 13.* 16 F 1911.

Hood, S. C., & True, R. H. Camphor cultivation in the United States. Yearbook U. S. Dept. Agr. **1910**: 449–460. *pl. 46–48.* 1911.

Jepson, W. L., Betts, H. S., & Mell, C. D. California tanbark oak. U. S. Forest Serv. Bull. 75: 1–34. *pl. 1–10.* 20 S 1911.

Johnson, D. S. Studies in the development of the *Piperaceae*—I. The suppression and extension of sporogenous tissue in the flower of *Piper Betel* L. var. *monoicum* C. DC. Jour. Exp. Zoölogy **9**: 715–749. *f. 1–71.*

Johnson, E. C. Timothy rust in the Unites States. U. S. Dept. Agr. Plant Ind. Bull. 224: 1–20. 4 Au 1911.

Jolivette, H. D. M. Spore formation in *Geoglossum glabrum* Pers. Trans. Wisconsin Acad. **16**: 1171–1190. *pl. 85–87.* 1910.

Jones, D. H. *Scolytus rugulosus* as an agent in the spread of bacterial blight in pear trees. Phytopathology **1**: 155–158. *pl. 23, 24.* O 1911.

Kellerman, K. F. Nitrogen-gathering plants. Yearbook U. S. Dept. Agr. **1910**: 213–218. *pl. 7–14.* 1911.

Kern, F. D. A biologic and taxonomic study of the genus *Gymnosporangium.* Bull. N. Y. Bot. Gard. **7**: 391–483. *pl. 151–161* + *f. 1–36.* 12 O 1911.

Includes descriptions of *Gymnosporangium fraternum, G. juvenescens,* and *G. effusum* spp. nov.

Kern, F. D. Two submerged species of *Uromyces.* Torreya **11**: 211–214. 18 O 1911.

Includes descriptions of *Uromyces seditiosus* and *U. argutus* spp. nov.

Kimball, L. F. Ferns of San Diego County, California. Fern Bull. **19**: 42–46. [O] 1911.

Kneucker, A. Bemerkungen zu den "Carices exsiccatae." Allgem. Bot. Zeits. **17**: 90–92. Je 1911; **17**: 118–121. Au 1911.

Kränzlin, F. *Orchidaceae-Monandrae-Dendrobiinae.* Pflanzenreich $4^{50\ II\ B21}$: 1–382. *f. 1–35.* 15 N 1910.

Léveillé, H. *Epilobium Arechavaletae* Lévl. nov. sp. Bull. Geog. Bot. **21**: 149. 1 S 1911.

Lignier, O. Notes anatomiques sur l'ovaire de quelques Papaveracées. Bull. Soc. Bot. France **58**: 279–284. *f. 1–3.* 21 Jl 1911; **58**: 337–344. *f. 4–8.* 21 Jl 1911; **58**: 429–435. *f. 9.* 5 S 1911.

Lipman, C. B. Nitrogen fixation by yeasts and other fungi. Jour. Biol. Chem. **10**: 169–182. O 1911.

McCulloch, L. A spot disease of cauliflower. U. S. Dept. Agr. Plant Ind. Bull. 225: 1–15. *pl. 1–3.* 29 Au 1911.

MacDougal, D. T. An attempted analysis of parasitism. Bot. Gaz. **52**: 249–260. *f. 1–6.* 17 O 1911.

Malaga Santolalla, F. Monografía del Departamento de Cajamarca 1906. Bol. Soc. Geog. Lima **20**: i–xiv + 1–326. 1906. [Illust.]

Marshall, R. The vegetation of Twin Island. Trans. Wisconsin Acad. **16**: 773–797. *pl. 62, 63.* [1910.]

Meyer, R. Über *Echinopsis Huatti* Lab., *Echinopsis apiculata* Lke., *Echinopsis Saliniana* Web. (hort. germ.) and *Echinopsis Bridgesii* S.-D. Monats. Kakteenk. **21**: 135–137. 15 S 1911.

Nash, G. V. The funkias or day-lilies. Jour. Hort. Soc. New York **1**: 164–171. O 1911.

Nieuwland, J. A. Some Linnaean trivial names. Am. Mid. Nat. **2**: 97–122. 15 S 1911.

Olive, E. W. Origin of heteroecism in the rusts. Phytopathology **1**: 139–149. O 1911.

Olsson-Seffer, P. Rubber and coffee in the Orient and Mexico. Am. Rev. Trop. Agric. **1**: 227–240. S 1910.

[Orcutt, C. R.] Flowerless plants of the United States. West Am. Sci. **18**: 9–40. S 1911.

Parmentier, P. Recherches anatomiques et taxonomiques sur les Juglandacées. Rev. Gén. Bot. **23**: 341–364. *pl. 8–11+f. 1.* 15 Au 1911.

Pax, F. *Euphorbiaceae-Adrianeae.* Pflanzenreich $4^{147\,II}$: 1–111. *f. 1–35.* 4 O 1910.

Pax, F. *Euphorbiaceae-Jatropheae.* Pflanzenreich 4^{147}: 1–148. *f. 1–45.* 3 My 1910.

Plummer, F. G. Chaparral. U. S. Forest Serv. Bull. 85: 1–48. *pl. 1–7+f. 1–7.* 17 Jl 1911.

Pretz, H. W. Flora of Lehigh County, Pennsylvania—I. Bull. Torrey Club **38**: 45–78. 7 Mr 1911. [Illust.]

Pritchard, F. J. The wintering of *Puccinia graminis tritici* E. & H. and the infection of wheat thru the seed. Phytopathology **1**: 150–154. *pl. 22+f. 1, 2.* O 1911.

Quehl, L. Bemerkungen über einige Arten von Mamillarien aus der Untergattung *Coryphantha* Engelm., Reihe *Aulacothelae* Lem. Monats. Kakteenk. **21**: 138–139. 15 S 1911.

Quehl, L. Kleine Mitteilungen und Fingerzeige. Monats. Kakteenk. **21**: 144. 15 S 1911.
Notes on *Echinocactus Fobeanus* Mieckley and *E. Pepinianus* Lem.

Reed, H. S. The effect of the club root disease upon the ash constituents of the cabbage root. Phytopathology **1**: 159–163. O 1911.

[Ridley, H. N.] *Castilloa* in Mexico. Agric. Bull. Straits and Fed. Malay States **10**: 212, 213. Jl 1911.

[Ridley, H. N.] *Cryptostegia grandiflora* in Mexico. Agric. Bull. Straits and Fed. Malay States **10**: 208, 209. Jl 1911.

Robinson, C. B. *Urticaceae* from the Sarawak Museum. Philip. Jour. Sci. **6**: (Bot.) 291–298. S 1911.

Scott, W. M., & Rorer, J. B. Apple blotch, a serious disease of southern orchards. U. S. Dept. Agr. Plant Ind. Bull. 144: 7–28. *pl. 1–6.* 6 Mr 1909.

Shimek, B. The prairies. Bull. Lab. Nat. Hist. Univ. Iowa 6^2: 169–240. *pl. 1–14.* O 1911.

Smith, E. F. Crown-gall and sarcoma. U. S. Dept. Agr. Plant Ind. Circ. 85: 1–4. 20 Je 1911.

Smyth, B. B., & Smyth, L. C. K. Catalogue of the flora of Kansas, Part I. Trans. Kansas Acad. Sci. **23, 24**: 273–295. Ja 1911.

Spaulding, P. The blister rust of white pine. U. S. Dept. Agr. Plant Ind. Bull. 206: 1–88. *pl. 1, 2+f. 1–5.* 22 Jl 1911.

Spaulding, P. European currant rust on the white pine in America. U. S. Dept. Agr. Plant Ind. Circ. 38: 1–4. 5 Au 1909.

Spaulding, P. The present status of the white-pine blights. U. S. Dept. Agr. Plant Ind. Circ. 35: 1–12. 23 Au 1909.

Spaulding, P. The timber rot caused by *Lenzites sepiaria.* U. S. Dept. Agr. Plant Ind. Bull. 214: 1–46. *pl. 1–4+f. 1–3.* 21 Jl 1911.

Spragg, F. A. Forage conditions of central Montana. Montana Agric. Exp. Sta. **36**: 5–40. *f. 1–23.* Je 1902.

Contains a generic key to the grasses and grasslike plants of Montana and an annotated list of the grasses of central Montana.

Stapf, O. *Aphelandra Fascinator.* Curt. Bot. Mag. IV. **7**: *pl. 8398.* O 1911.

A tropical South American species.

Sterrett, W. D. Scrub pine (*Pinus virginiana*). U. S. Forest Serv. Bull. 94: 1–27. *f. 1, 2.* 22 Jl 1911.

Sudworth, G. B., & Mell, C. D. "Colombian Mahogany." U. S. Forest Serv. Circ. 185: 1–16. *f. 1–11.* 3 Au 1911.

Sudworth, G. B., & Mell, C. D. Fustic wood: its substitutes and adulterants. U. S. Forest Serv. Circ. 184: 1–14. *f. 1–4.* 9 Je 1911.

Türckheim, H. von. Botanische Forschungsreise in Santo Domingo in den Jahren 1909 und 1910. Allgem. Bot. Zeits. **17**: 101–106. Au 1911.

Wadmond, S. C. Flora of Racine and Kenosha counties, Wisconsin: a list of the fern and seed plants growing without cultivation. Trans. Wisconsin Acad. **16**: 798–888. [1910.]

Walter, H. *Phytolaccaceae.* Pflanzenreich 4^{83}: 1–154. *f. 1–42.* 1 Je 1909.

INDEX TO VOLUME

New names and the final members of new combinations are in **heavier type**

Acacia, 475, 476, 480
Acaciaphyllites, 420
Acarospora sp., 293
Acer californicum, 336; rubrum, 225, 227–229, 231
Achillea lanulosa, 237, 239
Aconitum columbianum, 237; Lycoctonum, 313
Acrostichum citrifolium, 173
Actinella, 16; lanata, 239
Adiantum pedatum, 54
Adopogon, 12, 16
Adoxa moschatalina, 237
Aecidium alliicola, 375
Aeodes nitidissima, 511, 512
Agoseris, 16; acuminata, 17; attenuata, 17; altissima, 17; agrestis, 17; arachnoidea, 18, 21; aurantiaca, 18, 20, 237; elongata, 19; glauca, 17, 18, 21; gracilens, 20; gracilenta, 19; gracilenta Greenei, 19; graminifolia, 19, 20; grandiflora, 20, 21; Greeneana, 19; Greenei, 18, 19; humilis, 19; laciniata, 18, 21; Leontodon, 18; maculata, 17; major, 19; **obtusifolia,** 20; parviflora, 20; pubescens, 17; pumila, 17; roseata, 17; rostrata, 19, 20; scorzoneraefolia, 17; **tenuifolia,** 20; **turbinata,** 20; villosa, 17, 18
Agrimonia Bicknellii, 450; gryposepala, 449; hirsuta, 449; mollis **Bicknellii**, 450
Agropyrum Scribneri, 237
Alcea americana, 94; indica, 94
Alectoria bicolor, 291; divergens, 291; jubata implexa, 291; ochroleuca, 292; sarmentosa, 292
Algae of Lower California, Mexico, Some marine, 489
Algites, 420; **americana,** 401
ALLARD, H. A. Some common species of Crataegus at Thompson's Mills, Georgia, 25
Allium, 374; acuminatum, 374; brevistylum, 373, 374; pikeanum, 237, 241; reticulatum, 373; validum, 374
Alnus rhombifolia, 335, 336, 341, 342
Alsine baicalensis, 239; crassifolia, 239, 240
Alsinopsis obtusiloba, 237; propinqua, 237
Alsophila capensis, 546; decussata, 548, 550
Alternaria, 135, 137
Althaea, 94
Amarella plebeja, 238, 241
Amelanchier alnifolia, 335, 337; canadensis, 453; **nantucketense,** 453–455; oblongifolia, 453, 455; oblongifolia micropetala, 454; rotundifolia, 455; sanguinea, 455; spicata, 454, 455
Amnicola Yatesiana, 331
Amphicosmia multiflora, 547, 548
Amygdalaceae, 337
Amygdalus Persica, 460
Ananthacorus, 158, 159, 171–173, 190; angustifolius, 172, 176, 181, 188
Anatheca **dichotoma,** 502, 503, 513; furcata, 503
Anchistea virginica, 55
ANDREWS, F. M. Conjugation of two different species of Spirogyra, 299
Andromeda, 420, 421, 423; grandifolia, 420; Novae-caesareae, 420; Parlatorii, 420
Andropogon virginicus, 225, 231
Androsace carinata, 237; subumbellata, 237
Androvettia, 404
Anetium, 153, 156–159, 171, 173; citrifolium, 157, 173, 188
Angelica Grayi, 237
Anodonta cygnea impura, 331
Antennaria, 382; aprica, 237; media, 237, 241; microphylla, 237, 239; nardina, 237; rosea, 237, 239
Anthoceros levis, 221
Anticlea elegans, 239
Antrophyopsis, 175; immersum, 176
Antrophyum, 158, 159, 164–183, 187–190; anetioides, 169; Boryanum, 172; discoideum, 169; Dussianum, 169; ensiforme, 169; Jenmani, 169; latifolium, 156; lineatum, 171; minimum, 164; nanum, 175, 179, 188; plantagineum, 172, 174, 189; reticulatum, 169, 172, 176, 177, 181, 182, 190; semicostatum, 174, 175, 188; stenophyllum, 189; Werckleanum, 164; Williamsi, 190
Aphanolejeunea, 251, 254, 255, 271–273, gemmae of, 279–286; **crenata,** 276, 277, 286; **exigua,** 273–277, gemmae of, 281–285; **microscopica,** 273, 275, gemmae of, 281–284; **sicaefolia,** 277–279, 286
Aquilegia coerulea, 237; saximontana, 237
Arabis Drummondii, 237, 239
Aralia, 415; anisiloba, 417; groen-

landica, 408; spinosa, 516; washingtoniana, 409; Wellingtoniana, 415
Araliopsis, 413; **breviloba,** 417; **cretacea,** 413, 417; cretacea **dentata,** 416; cretacea **salisburiaefolia,** 416
Araucaria, 405, 420, 421; bladenensis, 405, 420; Jeffreyi, 420
Araucarites ovatus, 405
Arbutus, Dioecism in the trailing, with notes on the morphology of the seed, 531
Arbutus Menziesii, 335, 339, 342
Archilejeunea Sellowiana, 219; viridissima, 219
Arctostaphylos, 341; Manzanita, 335, 339, 340
Arenaria Fendleri, 237; sajanensis, 237
Argentina, 351; Anserina, 352–354, 447; argentea, 352, 354; Babcockiana, 353–355; Egedii, 353–356; litoralis, 352–355, 447, 448; occidentalis, 352–354; pacifica, 353; subarctica, 352, 353, 355
Aristida stricta, 231
Aronia arbutifolia, 452; atropurpurea, 452; nigra, 452
Artemisia Pattersonii, 237; scopulorum, 237
Arthonia patellulata alnicola, 287
Arthoniaceae, 287
ARTHUR, J. C. New species of Uredineae—VIII, 369
Arundinaria macrosperma, 227; tecta, 225, 226
Arundo, 422; groenlandica, 420
Asclepiodora viridis, 232
Aspergillus niger, 135, 137
Aspidiophyllum trilobatum, 410
Asplenium **cecilensis,** 403, 424; Dicksonianum, 409; ebenoides, 55; montanum, 56; pinnatifidum, 77; platyneuron, 57; platyneuron Hortonae, 58; Ruta-muraria, 57; Trichomanes, 58
Assimilation of atmospheric nitrogen by fungi, Upon, 135
Aster, 437; carolinianus, 232
Astrodapsis Whitneyi, 340
Athyrium acrostichoides, 58; angustifolium, 77; Filix-femina, 59
Aureae, 79

Baccharis, 229; glomeruliflora, 372; halimifolia, 229; oaxacana, 371; sordescens, 371; thesioides, 370
Baeomyces, 287; aeruginosus, 288
Baptisia perfoliata, 234; villosa, 232
BARTLETT, H. H. A supplemental description of Euphorbia ephedromorpha, 343
Batodendron arboreum, 516
BENEDICT, R. C. The genera of the fern tribe Vittarieae: their external morphology, venation, and relationships, 153
Benzoin melissaefolium, 232
Besseya alpina, 239
BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain—VII, 399
Betulaceae, 336
Biatora cinnabarina, 288; franciscana, 288; sphaeroides, 288
BICKNELL, E. P. The ferns and flowering plants of Nantucket, 103, 447
Biflorae, 80
Bistorta bistortoides, 237; 239, vivipara, 237, 239
Blackberries, 103; half high, 426; high, 426
BLANCHARD, W. H. Rubus of eastern North America, 425
Blastenia ferruginea, 290
Botrychium dissectum, 59; lanceolatum, 59; neglectum, 59; obliquum, 60, 76; simplex, 60; tenebrosum, 61; virginianum, 61
Boykinia Jamesii, 237
Brachiolejeunea bahamensis, 205, 218; corticalis, 218
Brachyphyllum macrocarpum, 422
BRAINERD, E. Further notes on the stemless violets of the South, 1; The caulescent violets of the southeastern United States, 191
Brassica, 478
Brevifoliae, 81
Brunnichia cirrhosa, 233
Bryum coronatum, 34
Buellia papillata, 287; scabrosa, 287

Caesalpinia, 423
Calamagrostis, 319; purpurascens, 237
Callicostella pallida, 36
Callymenia Requienii, 511
Caltha leptosepala, 237, 239, 240
Calycites, 420; diospyriformis, 418
Campanula rotundifolia, 237; uniflora, 237, 239
Camptosorus rhizophyllus, 61
Candicantes, 82
Canna flaccida, 232
Cannabinus, 94
Carduus scopulorum, 237, 239
Carex 422, 482; alpina, 237; capillaris, 237; chalciolepis, 237; Clarkii, 420; ebenea, 237; festiva, 237; lupulina, 481; lupulina **albomarginata,** 482; melanocephala, 237; petasata, 237; rupestris, 237; scopulorum, 237; Walteriana, 226
Carex lupulina, A new variety of, 481
Carinifex Sanctae-clarae, 331
Carnegiea, 473, 475, 476, 477; gigantea, 474

Castilleja lancifolia, 237; occidentalis, 237; sulphurea, 237, 239
Caudalejeunea Lehmanniana, 219
Caulinites, 402; inquirendus, 401
Caulescent violets of the southeastern United States, The, 191; synopsis of, 197
Celastrus, 537; arctica, 407; scandens, 537
Centroceras clavulatum, 492, 509; cryptacanthum, 509; cryptacanthum longiarticulatum, 509; oxyacanthum, 509
Celastrophyllum, 421, 423; crenatum, 420; elegans, 420
Cephalanthus occidentalis, 335, 339
Cephalotaxospermum carolinianum, 420
Ceramium clavulatum, 509
Cerastium beeringianum, 237; oreophilum, 237
Ceratiola ericoides, 225, 234
Ceratolejeunea cubensis, 213, 215; **integrifolia,** 213, 215, 222; patentissima, 214
Cercocarpus, 341; betulaefolius, 335, 337
Cetraria arctica, 292; cucullata, 292; islandica, 292; juniperina pinastri, 292; nivalis, 292; Oakesiana, 292; Richardsonii, 292
Chamaecyparis thyoides, 225, 235
Chamaenerion angustifolium, 237, 239, 241
Chamaerhodos, 367; erecta, 367; erecta parviflora, 367; Nuttallii, 367
Cheilanthes lanosa, 62
Cheilolejeunea decidua, 212; phyllobola, 210, 212; versifolia, 208
Chionophila Jamesii, 237
Chlorophyceae, 490
Cholisma ferruginea, 232, 516
Chondria obtusa paniculata, 508
Chondrophylla americana, 237, 239
Cinnamomum, 420, 421, 423; intermedium, 423; **Newberryi,** 420, 423
Cissites, 415, 417; Heerii, 415; Harkerianus, 414, 415; obtusum, 416; salisburiaefolius, 414–417
Cissus laciniata, 477
Citrophyllum, 423; aligerum, 420
Cladium effusum, 231, 235
Cladonia, 287; belliflora coccocephala, 288; cariosa, 289; cariosa Dawsoniana, 289; cariosa majuscula, 289; coccifera, 288; coccifera pleurota, 288; crispata, 288; crispata subcrispata, 288; digitata, 288; furcata scabriuscula, 288; pyxidata neglecta, 288; rangiferina, 288; rangiformis, 288; sp., 292; uncinalis, 288
Cladoniaceae, 288
Cladophlebis, 403, 409; Browniana, 409; **socialis,** 409
Cladophora Hutchinsiae, 491; **MacDougalii,** 491, 492, 513; ovoidea, 491, 492; trichotoma, 492
Claytonia megarrhiza, 237
Clementsia rhodantha, 237
Clethra alnifolia, 229
Cliftonia monophylla, 231, 233
Codium, 494; **decorticatum,** 494, 495; elongatum, 494, 495; galeatum, 493; Lindenbergii, 494; tomentosum, 493–495; tomentosum proliferum, 494
Coilodesme-?, 496; californica, 496
Collema melaenum, 290; melaenum polycarpum, 290
Collemaceae, 290
Cololejeunea, 251–261, gemmae of, 279–286; Biddlecomiae, 254, 256, gemmae of, 280, 281; calcarea, 251, 252, 254, 256, gemmae of, 280, 281; ceatocarpa, 265; ciliatilobula, 254; cuneifolia, 253; diaphana, 259, gemmae of, 282, 284, 285; erigens, 263; Goebelii, 265; heteromorpha, 257; Jooriana, 207, 270; lanciloba, 253, 268; longistylis, 282; Macounii, gemmae of, 281–283; marginata, 262; microscopica, 273; minutissima, 258, 259, gemmae of, 282, 283, 285; **myriocarpa,** 256, 258, 259, gemmae of, 281–283, 285; ovalifolia, 281; scabriflora, 263
Colpomenia tuberculata, 495; sinuosa tuberculata, 495
Colutea obovata, 407; primordialis, 407
Comarum, 81, 356; angustifolium, 357; palustre, 356; tomentosum, 357
Conferva trichotoma, 492
Conioselinum scopulorum, 237, 239
Conjugation of two different species of Spirogyra, 299
Contributions to the history and bibliography of the roselle, 91
Contributions to the Mesozoic flora of the Atlantic coastal plain—VII, 399
Cordaites, 406
Cornaceae, 338
Corneocyclas compressa, 331
Cornus, 408; **cecilensis,** 408, 424; florida, 516; glabrata, 335, 338, 342
Corydalis nobilis, 312; ochroleuca, 313
Crataegus, 25–32, 437–458; Beadlei, 25, 29–32; **Bicknellii,** 243, 244, 455–459; chrysocarpa, 244, 455, 457; collina, 25, 29, 32; Crus-galli, 25, 26, 32, 456; Jonesae, 243; macrosperma, 458; pruinosa, 455, 457, 458; rotundifolia, 457; rotundifolia Bicknellii, 244; **schizophylla,** 243, 455; spathulata, 25, 27, 32; tomentosa punctata, 459; uniflora, 25, 26, 28; Williamsii, 243
Credneria rhomboidea, 413
Crepidispermum, 16
Crepis, 12, 15; acuminata, 14; alpicola, 13; ambigua, 15, 16; angustata, 13; barbigera, 14; biennis, 13; denticulata, 13; exilis, 14; glauca, 12, 13; glaucella,

12, 13; gracilis, 13, 14; intermedia, 13; occidentalis, 14; occidentalis costata, 14; occidentalis gracilis, 13; perplexans, 12, 13; petiolata, 12, 13; platyphylla, 12, 13; pumila, 14; riparia, 12, 13; riparia parva, 13; runcinata, 12; runcinata alpicola, 13; runcinata hispidulosa, 13; scopulorum, 13; scopulosa, 13; **seselifolia,** 14; tomentulosa, 12, 13
Crossotolejeunea, 252
Croton, 423
Crotonophyllum, 420, 423
Cryptomeric inheritance in Onagra, 461
Cunninghamites elegans, 422
Cyathea, 545; arborea, 550; capensis, 545; horrida, 545; multiflora, 545, 547, 548, 550
Cyathea multiflora, type of the genus Hemitelia R. Br., On the identity of, 545
Cyclodictyon Liebmanni, 36
Cyclolejeunea, gemmae of, 279; accedens, 274
Cypripedium Reginae, 232

Dalbergia, 407; **severnensis,** 407, 424
Dammara borealis, 400; cliffwoodensis, 400
Dasiphora, 356, 362; fruticosa, 237, 238
Dasystephana Parryi, 238; Romanzovii, 238
Datura, 541; laevis, 541
Delphinium, 313; azureum, 308; decorum scaposum, 307; nudicaule, 308; scaposum, 307–310, 312, 317
Dennstaedtia punctilobula, 62
Dentaria diphylla, 192
Deschampsia alpicola, 237; caespitosa alpina, 237
Description of two new fossil figs from Wyoming and Montana, 389
Development of the embryo sac and endosperm in some seedless persimmons, 379
Dewalquea, 422; Smithi, 420
Dewberries, 426
Dicranella Hilariana, 33; Liebmanniana, 33; mexicana, 33; Perrotettii, 33; tenuirostris, 33; trematodontifolia, 33
Dictyota Bartayresiana, 498; Binghamiae, 499; crenulata, 499; **Vivesii,** 497–499, 512
Dioecism in the trailing arbutus, with notes on the morphology of the seed, 531
Diospyros, 418, 421, 423; brachysepala, 418; primaeva, 417, 418, 420; rotundifolia, 420; **vera,** 418; virginiana, 418, 379; virginiana Turneri, 339
Diplasiolejeunea, 255, 262, 267; pellucida, 262; Rudolphiana, 207; unidentata, 207
Dodecatheon pauciflorum, 237, 239
Dodonaea, 295
Doryanthites, 406; **cretacea,** 406
Draba aurea, 238; Parryi, 238; streptocarpa, 238
Drepanolejeunea Araucariae, 276; infundibulata, 274
Dryas octopetala, 238
Drymocallis, 80, 86, 362; agrimonioides, 362–367; albida, 362, 365; amplifolia, 362, 363; arguta, 362, 366, 367; arizonica, 362, 363; ashlandica, 366, 367; convallaria, 362, 366, 367; corymbosa, 362; cuneifolia, 366; fissa, 363, 366, 367; foliosa, 362, 363; glabrata, 363, 366, 367; glandulosa, 363, 364, 367; glandulosa incisa, 362; glandulosa monticola, 362; glutinosa, 362, 366, 367; gracilis, 366; Hanseni, 363, 366; incisa, 366; lactea, 366; laxiflora, 362, 364, 365; micropetala, 362, 365; oregana, 362, 363, 365; pseudorupestris, 366, 367; pumila, 362, 363; reflexa, 364–366; rhomboidea, 363, 366, 367; valida, 362, 363, 367; viscida, 365, 367; viscosa, 362, 363; Wrangelliana, 363–367
Dryopteris Boottii, 63; Clintoniana, 77; cristata, 63, 64; cristata × intermedia, 63; cristata × marginalis, 64; Goldiana, 65, 76; intermedia, 65, 66; intermedia × marginalis, 65; marginalis, 66; noveboracensis, 66; simulata, 77; spinulosa, 66; Thelypteris, 66
Duchesnea, 357; indica, 357

Early spring aspects of the coastal plain vegetation of South Carolina, Georgia, and northeastern Florida, 223
Echinocactus, 477
EGGLESTON, W. W. New Crataegi of the northeastern manual range, 243
Eichhornia crassipes, 382
Elephantella groenlandica, 238, 239
Enteromorpha compressa, 491; intestinalis, 490; Linza, 490
Ephebe pubescens, 290
Epidendrum conopseum, 516
Epigaea, 531–542; dioecism in, 535; evidence of heterostyly in, 534; morphology of the seed, 538; repens, 531, 533, 537, 538, 540, 541, 543
Epilobium anagallidifolium, 238; ovatifolium, 238; stramineum, 238, 239
Equisetum arvense, 73; fluviatile, 73; hyemale, 73; hyemale affine, 74; sylvaticum, 74
Erianthus sp., 225, 226
Ericaceae, 339
Ericoideae, 540
Erigeron elatior, 238; formosissimus, 238, 239; leucotrichus, 241; pinnatisectus, 238; simplex, 238, 239, 241; Smithii, 238, 239; uniflorus, 241

Eriocaulon decangulare, 229
Eritrichium argenteum, 238
Erysimum oblanceolatum, 238, 239
Euantrophyum, 175, 176
Eucalyptus, 334, 421, 423; angusta, 421; Geinitzi, 421; Wardiana, 421
Eulejeunea albida, 208; flava, 207
Euonymus europaeus, 537
Euosmolejeunea clausa, 212; duriuscula, 212; opaca, 212; trifaria, 213
Eupatorium rotundifolium, 229
Euphorbia ephedromorpha, 343–346
Eupolytaenium, 171
Eusargassum, 496
Euvittaria, 166
EVANS, A. W. Hepaticae of Puerto Rico, 251; The Hepaticae of the Bahama Islands, 205
Evernia prunastri thamnodes, 292
Evidence of recapitulation in the Vittarieae, 186

Fabaceae, 338
Fagaceae, 336
Fauchea, 507; laciniata, 506; **mollis**, 507, 508, 513; nitophylloides, 507, 508; repens, 506; **Sefferi,** 506, 513
Ferns and flowering plants of Nantucket, The, 103, 447
Festuca brachyphylla, 238; ovina ingrata, 238
Ficus, 389–392, 400, 420; atavina, 421; Bruesi, 391; callicarpa, 391; **Ceratops**, 389; crassipes, 421; elastica, 400; Krausiana, 421; **matawanensis**, 399, 424; neurocarpa, 391; ovata, 410; ovatifolia, 410; protogaea, 391; punctata, 391; reticulata, 400; **Russelli,** 392; Stephensoni, 421; Woolsoni, 410
Figs from Wyoming and Montana, Description of two new fossil, 389
Filix bulbifera, 67; fragilis, 67, 238
Fissidens polypodioides, 33
Flabellaria, 405
Flora of Lehigh County, Pennsylvania, 45
Fragaria, 357; americana, 358, 359, 361; bracteata, 358; californica, 357; canadensis, 358; chiloensis, 360, 361; cuneifolia, 360; firma, 359; glauca, 238–240, 359, 360; grandiflora, 357, 360, 361; Grayana, 360, 361; illinoensis, 360; insularis, 359; mexicana, 357; multicipita, 357, 358; ovalis, 359; pauciflora, 359; platypetala, 358, 359; prolifica, 359; pumila, 359; Suksdorfii, 359, 360; Terrae-novae, 358; vesca, 357–361, 447; virginiana, 357–361, 447; yukonensis, 359, 360
Frasera carolinensis, 232
Fraxinus oregona, 336
Frigidae, 79
Frullania, 205, 256; arietina, 220; caroliniana, 220; gibbosa, 220; obcordata, 220; riojaneirensis, 220; squarrosa, 220
Fucus tomentosus, 493–495
Further notes on the stemless violets of the South, 1
Fusarium, 135, 137; Zeae, 135

GAGER, C. S. Cryptomeric inheritance in Onagra, 461
Gelsemium sempervirens, 516
Gemmae of Cololejeunea, Leptocolea, and Aphanolejeunea, 279
Genera of the fern tribe Vittarieae: their external morphology, venation, and relationships, The, 153
Gentiana cruciata, 312, 313; Parryi, 238; plebeja, 238, 239, 241; Romanzovii, 238
Georgia, Some common species of Crataegus at Thompson's Mills, 25
Geum agrimonioides, 362; canadense, 449; virginianum, 449
Gigartina Chamissoi, 502; Chauvinii, 502; papillata, 502; Teedii, 502
Ginnania undulata, 502
Glandulosa, 80
Gleichenia gracilis, 402; Saundersii, 402, 424
Glossophora Kunthii, 499
Gonidea angulata Cooperi, 331
Gracilaria, 504, 508; armata, 504; confervoides, 504; compressa, 504; Cunninghamii, 504; multipartita, 504; peruana, 504, 505; **Vivesii,** 503–505, 513
Graciles, 83; pinnatae, 83, 85
Grammitis pumila, 162
Grossularia Menziesii, 335, 337
Grossulariaceae, 337
Gymnogramma, 163

Halimeda, 493; discoidea, 492, 493; Tuna, 493
Halymenia, 510; **actinophysa,** 509–513; coccinea, 511; floridana, 510, 511; latifolia, 510; ulvoidea, 510, 511
Hamamelites, 423; cordatus, 421
HANNIBAL, H. A Pliocene flora from the Coast Ranges of California, 329
HARPER, R. M. Early spring aspects of the coastal plain vegetation of South Carolina, Georgia, and northeastern Florida, 223; The relation of climax vegetation to islands and peninsulas, 515
HARRIS, J. A. Teratological fruits of Ptelea, 385
Hecistopteris, 156, 157, 159, 162, 179, 182, 189; pumila, 162; Werckleana, 155, 164
Heckeria umbellata, 382
Hedera, 423; primordialis, 421
Hemionitis brasiliana, 169; cayennensis,

169; lanceolata, 169; lineata, 169; reticulata, 174
Hemitelia, 545; Hartii, 548; multiflora, 546, 548, 550; nigricans, 547, 548, 550; obscura, 548, 550; Parkeri, 546
Hemitelia R. Br., On the identity of Cyathea multiflora, type of the genus, 545
Hepaticae of Puerto Rico, 251
Hepaticae of the Bahama Islands, The, 205
Heteranthera limosa, 382
Heterolepis, 420
Heteropleura, 16; ambigua, 15, 16; **Fendleri,** 15, 16
Heterothecium sanguinarium, 288
Heuchera Hallii, 238; parvifolia, 238
Hibiscus, 97; digitatus, 97; gossypifolius, 97; Sabdariffa, 91, 97
Hicoria, sp., 516
Hieracium, 14, 15; albertianum, 15; albiflorum, 15; canadense, 14, 15; columbianum, 14, 15; cynoglossoides, 15; Fendleri, 15; griseum, 15; macranthum, 15; nigrocollinum, 15; runcinatum, 12, 13; scabriusculum, 15; Scouleri, 15; umbellatum, 14, 15
Holomitrium crispulum, 33
Hordeum jubatum, 377, 378
HOWE, M. A. Phycological studies—V. Some marine algae of Lower California, Mexico, 489
HOWE, R. H. JR. List of lichens collected in the Yukon region by Mr. R. S. Williams, 287
Hyophila Oerstediana, 34
Hypecoum Matthioli, 94
Hypopterygium Tamarisci, 36
Hyptis stellulata, 369
Ilex, 407; glabra, 226, 231; myrtifolia, 226, 231–233; opaca, 516; **severnensis,** 407, 424
Illicium, 420, 423
Indica elegans, 94
Induced and occasional parasitism, 473
Iris hexagona, 232
Isoetes, 154; canadensis, 76; Engelmanni, 76
Isopappus divaricatus, 225
Ivesia, 80

Jatropha stimulosa, 378
Juglans, 422; arctica, 420
Juncoides parviflorum, 238; spicatum, 238
Juncus castaneus, 238; Drummondii, 238; Roemerianus, 229; triglumis, 238
Jungermannia cuneata, 264; flava, 207; marginata, 262, 266; microscopica, 273
Jungermanniaceae, 206
Juniperus, 235; barbadensis, 229, 235; virginiana, 231, 235
Jussiaea grandiflora, 229

KINGSLEY, M. A. On the anomalous splitting of the rhizome and root of Delphinium scaposum, 307
KNOWLTON, F. H. Description of two new fossil figs from Wyoming and Montana, 389
Krameria, 479
Krigia, 12, 16; virginica, 12
Kuhnistera pinnata, 225, 234

Lactuca, 21, 22; augustana, 22, 23; integrifolia, 23; **polyphylla,** 23; saligna, 22; Scariola, 22; Scariola integrata, 22; virosa, 22, 23
LAMSON-SCRIBNER, F. Notes on certain species of Muhlenbergia, 319
Laurencia glandulifera, 508; paniculata, 508; patentiramea, 508; pinnatifida, 508
Laurophyllum, 421, 423; elegans, 420; nervillosum, 420
Laurus, 421, 423; atanensis, 420; canariensis, 339; plutonia, 420
Lecanora calcarea, 290; castanea, 290; epibrya, 290; lentigera, 289; pallida, 289; rugosa, 290; Schleicheri, 290; subfusca argentata, 289; subfusca campestris, 289; thamnoplaca, 289; ventosa, 290
Lecanoraceae, 289
Lecidaceae, 287
Lecidea alpestris, 293; arctica, 293; contigua hydrophila, 288; platycarpa, 288; sp., 293
Leguminosites, 420–423; robiniifolia, 420
Lehigh County, Pennsylvania, Flora of, 45
Lejeunea, 251–256, 261; americana, 207; calcarea, 252, 253; cardiocarpa, 267, 268; cavifolia, 252; ceatocarpa, 265; ciliatilobula, 254; cuneata, 264, 265; cuneifolia, 280; epiphyta, 257, 258; erigens, 262, 264; flava, 207, 208; glaucescens, 208; Goebelii, 265; heteromorpha, 257, 258; heterophylla, 273, 275; hyalino-marginata, 266; Jooriana, 270; laevigata, 253, 254; liliputiana, 279; Macounii, 281; marginata, 262, 266, 267; microscopica, 252, 253, 273; micrandroecia, 262; Moorei, 207; myriocarpa, 256–258; obtusangula, 215; obliqua, 254, 265; ovalifolia, gemmae of, 281–284; pseudostipulata, 256; scabriflora, 262, 264, 265; sicaefolia, 273, 277, 278; Sintenisii, 273, 275; Wrightii, 257
Lepidopilum brevipes, 36; polytrichoides, 36; semilaeve, 36
Leptasea austromontana, 238; chrysantha, 238; flagellaris, 238
Leptocolea, 251–255, 260–262, 267, gemmae of, 279–286; **cardiocarpa,** 260, 268–271, 286; **ceatocarpa,** 265; gemmae of, 282–284; **cuneifolia,** 253, 280–284,

gemmae of, 281, 282, 284; **Goebelii**, 265, 280, gemmae of, 282–285; **Jooriana**, 270, 271, gemmae of, 282, 283, 285; laevigata, 253; **lanciloba**, 253, 268, gemmae of, 282, 283; **longistylis**, 282, 283; marginata, 262, 266, 268; **micrandroecia**, 252, 262; **ovalifolia**, 281–284; **planifolia**, 265–268, 270, 286; **scabriflora**, 262, 265, gemmae of, 282–286
Leptogium myochroum saturninum, 290
Leptolejeunea elliptica, 218
Leucobryum giganteum, 33
Leucodoniopsis plicata, 34
Leucolejeunea xanthocarpa, 220; unciloba, 219
Leucophyllae, 82, 84
Leucothoe Catesbaei, 322
Lewisia pygmaea, 238, 239, 241
Lichens collected in the Yukon region by Mr. R. S. Williams, List of, 287
Lindigia aciculata, 35
Liquidambar Styraciflua, 225, 227, 516
Liriodendron Tulipifera, 225
List of lichens collected in the Yukon region by Mr. R. S. Williams, 287
List of plants collected above timber line on Pikes Peak, with altitudinal extensions and notes, 237
Lloydia serotina, 238
Lopholejeunea Sagraeana, 219
Lorinseria areolata, 77
Loxogramma, 183
Lubie Endigi, 95
Lupinus diffusus, 225, 234
Luzula parviflora, 238; spicata, 238
Lychnis montana, 238
Lycopodium, 420, 421; annotinum, 78; clavatum, 74; complanatum, 74, 78; complanatum flabelliforme, 74; inundatum, 75; lucidulum, 75; obscurum, 75; palmatum, 68; tristachyum, 78
Lygodium palmatum, 68

MacDougal, D. T. Induced and occasional parasitism, 473
Machaeranthera aspera, 238, 239
Mackensen, B. Two new species of Opuntia, 141
Macromitrium apiculatum, 34; **flavopilosum**, 34; mucronifolium, 34; ulophyllum, 34
Macrorhynchus glaucus laciniatus, 18, 21; purpureus, 18, 19
Magnolia, 421, 423; californica, 339; Capellinii, 406, 420; Fraseri, 232; glauca, 225–229, 231; grandiflora, 226–231, 234, 516; Newberryi, 420; obtusata, 420; tenuifolia, 420
Malaceae, 337
Malapoenna geniculata, 232
Malus Malus, 452
Malva, 94
Marchantia chenopoda, 206
Marchantiaceae, 206
Marshallia trinervia, 232
Mastigolejeunea auriculata, 219
Matteuccia Struthiopteris, 68
Mayepea **macrocarpa**, 145
Maxon, W. R. On the identity of Cyathea multiflora, type of the genus Hemitelia R. Br., 545
Melia Azederach, 225
Mertensia alpina, 238, 241; ciliata, 238, 241; picta, 241
Mesadenia sulcata, 232
Mesophytic climax, 517
Meteoriopsis patula, 35; recurvifolia, 35
Metzgeriopsis, 251, 252; pusilla, 251
Micranthes rhomboidea, 238, 240
Microlejeunea bullata, 208; laetevirens, 208; lucens, 208
Mitchella repens, 516
Momisia, 420, 422
Monogramma, 153, 156, 158–163, 167, 182, 190; dareicarpa, 160–163, 178, 180, 182, 185, 186, 188; graminea, 160–163, 166, 178, 182, 185, 187, 189; paradoxa, 160–162, 164, 167, 168, 178, 186, 189; subfalcata, 160, 161, 178, 179, 180, 186, 189; trichoidea, 160–162, 178, 186, 189
Monotropoideae, 540
Moriconia americana, 422
Morus alba, 145; celtidifolia, 146; **mollis**, 145
Muhlenbergia, Notes on certain species of, 319
Muhlenbergia californica, 323–325; comata, 319; **curtifolia**, 328; curtifolia **Griffithsii**, 328; glomerata brevifolia, 323; huachucana, 325; Lemmoni, 324–327; neo-mexicana, 320–322; Parishii, 323; pauciflora, 320–322; **polycaulis**, 327; Pringlei, 320; sp., 377; sylvatica californica, 323; sylvatica Pringlei, 320, 321
Multifidae, 82, 86
Multijugae, 80, 82, 83
Myrica, 421, 422; cerifera, 226, 234, 516; Brittoniana, 420; elegans, 420
Myrsine, 423; Gaudini, 408, 420

Nantucket, The ferns and flowering plants of, 103, 447
Narinam-Poulli, 96
Neckera disticha, 35; undulata, 35
Nematostylae, 80, 356
Nephroma arcticum, 291; expallidum, 291
New Crataegi of the northeastern manual range, 243
New species of Uredineae, 369
New variety of Carex lupulina, A, 481
Nitophyllum ocellatum, 508
Nitrogen, Upon assimilation of atmospheric, by fungi, 135

Notes on certain species of Muhlenbergia, 319
Notes on Rosaceae, 79, 351
Notes on Rutaceae—V. Species characters in Ptelea and Taravalia, 295
Nymphaea sagittifolia, 232
Nyssa biflora, 225–231; capitata, 232; Ogeche, 228, 231, 232; sylvatica, 226; uniflora, 228, 229, 233, 234

Octoblepharum albidum, 33
Olsynium grandiflorum, 376
Onagra, Cryptomeric inheritance in, 461
Onagra, 465; biennis, 461–463, 471; rubricalyx, 468; rubrinervis, 468
Onoclea, 402, 421; inquirenda, 401, 420, 424; sensibilis, 68
On the anomalous splitting of the rhizome and root of Delphinium scaposum, 307
On the identity of Cyathea multiflora, type of the genus Hemitelia R. Br., 545
Onychiopsis, 409
Ophioglossum vulgatum, 77
Opuntia, 474, 477, 478, 480; Blakeana, 474, 476, 478; discata, 475; **leptocarpa**, 141; Lindheimeri, 142; macrorhiza, 142, 143; **Roseana**, 142; versicolor, 477; vulgaris, 516
Oreobroma pygmaea, 238, 240
Oreochrysum Parryi, 238
Oreoxis humilis, 238, 239
Orontium aquaticum, 234
Orthostichopsis tetragona, 35
Osmanthus americana, 516
Osmunda, 402; cinnamomea, 69, 229; cinnamomea frondosa, 69; Claytoniana, 69; obergiana, 401, 402; spectabilis, 69
Oxyria digyna, 238

Pachistima, 420, 423
Pachypteris, 404
Padina Durvillaei, 497; sp., 497
Padus demissa, 335, 337
Palmocarpon sp., 389, 390
Paludestrina longinqua, 331
Panama mosses, 33
Pannaria brunnea, 291; lepidiota coralliphora, 291; pezizoides, 291
Pannariaceae, 291
Parkinsonia, 476
Parmelia conspersa, 291; stygia, 291
Parmeliaceae, 291
Paronychia pulvinata, 238
Pasania densiflora, 335, 336, 342
Pecopteris socialis, 409
Pedicularis Parryi, 238
Pellaea atropurpurea, 69
Peltigeraceae, 291
Penicillium, 135, 138; glaucum, 138
PENNINGTON, L. H. Upon assimilation of atmospheric nitrogen by fungi, 135
Pennsylvania, Flora of Lehigh County, 45
Pentstemon glaucus stenosepalus, 238, 239; Hallii, 238
Persea Borbonia, 516
Persimmons, Development of the embryo sac and endosperm in some seedless, 379
Pertusaria dactylina, 289; rhodocarpa, 289
Phacelia sericea, 238
Phaeophyceae, 495
Phalaris, 375
Phaseolus indicus, 95
Phegopteris Dryopteris, 77; hexagonoptera, 70; Phegopteris, 71
Philonotis sphaericarpa, 34; nanodendra, 34
Phleum alpinum, 238
Phoradendron flavescens, 225, 226–231, 516
Phragmites, 422; Pratti, 420
Phycological studies—V. Some marine algae of Lower California, Mexico, 489
Phyllogonium fulgens, 35; viscosum, 35
Physa heterostropha, 331
Physocolea, 252
Physostegia virginiana, 541
Pieris nitida, 231
Pilophorus cereolus acicularis, 289
Pilotrichum amazonum, 35
Pinaceae, 335
Pinus, 422; caribaea, 519, 520; echinata, 225–229; Elliottii, 226, 231, 233, 236, 520; glabra, 227, 232, 235; palustris, 225–231, 515, 519–524; raritanensis, 421; rigida, 524; serotina, 225, 226, 228, 231; Taeda, 225–228, 231
Piper medium, 382
Pistia Nordenskioldi, 405, 406
Placodium elegans, 290; ferrugineum, 290; Jungermanniae, 290; vitellinum, 290
Plagiochila ludoviciana, 206
Planorbis trivolvis, 331
Platanus, 412, 415; Heerii, 409, 411–413; recurvata, 411, 412, 415
Pleurogramma, 156, 165
Pliocene flora from the Coast Ranges of California, A, 329
Pluchea bifrons, 229
Poa arctica, 238; Grayana, 238; phoenicea, 238; rupicola, 238
Podeum, 504, 508
Podozamites Knowltoni, 403, 421; lanceolatus, 410; marginatus, 406, 410
Polemonium confertum, 238; pulcherrimum, 238
Polysiphonia californica, 508

Polystichum acrostichoides, 71
Polypodium marginellum, 157; polypodioides, 516; serrulatum, 157; vulgare, 71
Polypogon, 319
Polytaenium, 159, 169–190; **anetioides,** 169; **brasilianum,** 169; **cayennense,** 169, 174, 179, 189; **discoideum,** 169, 190; **Dussianum,** 169; **ensiforme,** 169, 171; **Jenmani,** 169, 171; **lanceolatum,** 169, 170, 172, 176, 177, 180, 181, 190; lanceolatum Feei, 189; lineatum, 169, 170, 172; **quadriseriatum,** 170–172, 178, 179, 190
Pontederia cordata, 229, 382
Populus trichocarpa, 335, 336
Porotrichum cobanense, 35
Potamogeton, 422
Porphyra leucosticta, 499
Potentilla, 79, 351; acuminata, 80; alpestris, 79; ambigens, 85; Anserina, 351, 352, 354; Anserina Egedii, 353; Anserina grandis, 352, 353; Anserina groenlandica, 353; Anserina sericea, 352; anserinoides, 354; arachnoidea, 89; argentea, 448; arguta, 362; arizonica, 86; argyrea, 82; atrovirens, 86, 89; biflora, 80; bipinnatifida, 86, 87, 238, 239; brevifolia, 80, 81; Breweri, 83; Bruceae, 82, 83; canadensis, 357, 448, 449; cascadensis, 85; ciliata, 367; coloradensis, 83; crinita, 84, 85; decurrens, 85; dissecta decurrens, 85; diversifolia pinnatisecta, 84, 85; diffusa, 83; Drummondii, 85; Egedii, 353; effusa, 82, 83; emarginata, 79; filipes, 83, 238; fissa, 367; flabelliformis, 79; fragiformis, 79; fragiformis parviflora, 79; Friesiana, 79; glabrella, 86, 87, 89; glandulosa, 364, 366, 367; glandulosa fissa, 366; glandulosa genuina, 366; glandulosa nevadensis, 367; glaucophylla, 238; glutinosa, 367; gracilis, 83; Hickmani, 84; Hippiana, 82, 83; Hippiana pulcherrima, 83; klamathensis, 83, 84; Keilhavii, 87; lactea, 366, 367; lasiodonta, 86; Lemmoni, 84, 85; litoralis, 86, 88; lupina, 82; lupulina, 82; Macounii, 81; maculata, 79; millefolia, 84; minutifolia, 82; missourica, 88, 89; monidensis, 85; monspeliensis, 448; multifida, 86, 88; nana, 79; Nelsoniana, 84, 85; nivea, 79; occidentalis, 352; oregana, 364; Osterhoutiana, 83; ovina, 84, 85; pacifica, 352–354; palustris, 80, 81, 356; palustris villosa, 357; paucijuga, 86; pectinata, 89; pennsylvanica, 82, 86–89; pennsylvanica communis, 88; pinnatifida, 86; pinnatisecta, 84, 85; plattensis, 84, 85, 89; propinqua, 83, 86; proxima, 81; pseudosericea, 86, 87; pulchella, 86, 87; pulchella elatior, 87, 88; pulcherrima, 83; pumila, 449; reflexa, 364, 367; reptans, 357; rhomboidea, 367; Richardii, 85; rosulata, 80; rubricaulis, 81, 82; rubripes, 81–83, 86; rupestris, 366; rupestris americana, 366; rupicola, 83; Salesoviana, 356; saximontana, 82, 238; saxosa, 80, 81; sericea glabrata, 89; simplex, 448; Sommerfeltii, 87; strigosa, 86, 89; subarctica, 88, 353; trichocarpa, 80, 356; tridentata, 80; Vahliana, 87; versicolor, 83, 84; virgulata, 86; viridior, 82, 83, 238, 239; Wahliana, 87; wyomingensis, 84, 85; Wrangelliana, 367
Prenanthes pauciflora, 12
PRETZ, H. W. Flora of Lehigh County, Pennsylvania, 45
Primula angustifolia, 238; Parryi, 238
Prionodon densus, 35
Prionolejeunea, 252
Protea, 422
Proteoides, 420–422; lancifolius, 421
Protophyllocladus, 420, 421; **lobatus,** 403; polymorphus, 404; subintegrifolius, 404, 405
Protophyllum multinerve, 411; Sternbergii, 411
Prunus americana, 460; angustifolia, 255; Avium, 459; Cerasus, 459; maritima, 459; serotina, 459
Pseudocryphaea flagellifera, 34
Pseudocymopterus multifidus, 238, 239; purpureus, 238, 239
Pseudotsuga taxifolia, 335
Psoralea physodes, 335, 338
Ptelea, 295–297; angustifolia, 295; Baldwinii, 295; monophylla, 234; tomentosa, 297; trifoliata, 295–297, 385; trifoliata mollis, 297; viscosa, 295
Ptelea, Teratological fruits of, 385
Pteridium aquilinum, 72
Pteridophyta, 54
Pteris, 403; Albertsii, 403; angustifolia, 172; graminea, 159; lineata, 164
Pterospermites, 411
Ptilocalais **macrolepis,** 11; major, 11; nutans, 11
Ptilophora major, 11
Ptiloria, 11; pauciflora, 12; ramosa, 11, 12
Puccinia Allii, 375; Asparagi, 375; Blasdalei, 374, 375; Calochorti, 376; **egregia,** 370; **exornata,** 370; **fidelis,** 369; granulispora, 375; mutabilis, 373, 375; oaxacana, 370; **pagana,** 372, 376; **pistorica,** 372; Porri, 375; rubigovera, 378; Sisyrinchii, 376; **sphenica,** 371; Winteriana, 375
Puerto Rico, Hepaticae of, 251
Pyroloideae, 540
Pyrus communis, 452

Quercus, 420, 422; agrifolia, 335, 337; Catesbaei, 225, 227, 231; Chapmani, 516; chrysolepis, 335–337, 342; falcata, 229; geminata, 516; laurifolia, 516; marylandica, 225; nigra, 225–229, 231; Phellos, 229; virginiana, 229, 516, 519

Radiovittaria, 166
Radula, 251; australis, 206
Ramalina polymorpha, 292; pusilla, 292; minuscula, 292
Ranunculaceae, 311
Ranunculoides, 79–81
Ranunculus inamoenus, 238, 239; multifidus, 382
Relation of climax vegetation to islands and peninsulas, The, 515
Rectolejeunea Berteroana, 208, 210; **Brittoniae,** 209, 211, 212, 222; flagelliformis, 210; phyllobola, 211, 212
Relationships of the Vittarieae, 183
Rhamnaceae, 338
Rhamnus californica, 335, 338, 342; Purshiana, 335, 338, 342
Rhizogonium spiniforme, 34
Rhodiola integrifolia, 238, 242; polygama, 238, 242
Rhodophyceae, 499
Rhodophyllis bifida, 508
Rhodymenia, 504; peruviana, 505
Rhodymenioideae, 508
Rhus, 423
Rhynchostegium scariosum, 36
Ribes lentum, 238, 241
Ricciaceae, 206
Ricciella crystallina, 206
Rivales, 85
Robinia hispida, 232
Roripa alpina, 238, 239
Rosa, 437; carolina, 450; cinnamomea, 451; humilis, 451; rubiginosa, 451; rugosa, 451; sp., 450; virginiana, 450
Rosaceae, 103, 337, 447
Rubiaceae, 339
Rubricaules, 81
Rubus of eastern North America, 425
Rubus, 425, 429, 432, 435–439; abbrevians, 119; allegheniensis, 106, 426–431, 435, 438; allegheniensis × argutus, 111; allegheniensis × frondosus, 112, 114; allegheniensis × procumbens, 113; amabilis, 436; amicalis, 436; Andrewsianus, 106, 426, 429, 430, 434, 438; angulatus, 439; arenicolus, 123; argutus, 106, 116, 118, 429; argutus × Baileyanus, 116; argutus × flagellaris, 117, 119; argutus × frondosus, 114; argutus × nigricans, 115; argutus × procumbens, 116, 121; ascendens, 115; Baileyanus, 109, 116, 119, 122, 128, 130, 131, 438; Baileyanus × Enslenii, 130; Baileyanus × flagellaris, 132; Baileyanus × frondosus, 118, 119, 121; Baileyanus × procumbens, 127; biformispinus, 120; Boyntoni, 438; caesius, 432; canadensis, 125, 426, 430, 432, 435; canadensis frondosus, 435; canadensis **Millspaughii,** 427; cuneifolius, 426, 434; Enslenii, 109, 123, 131, 132, 438; Enslenii × flagellaris, 131; Enslenii × frondosus, 123; Enslenii × hispidus, 125; Enslenii × procumbens, 128; flagellaris, 108–110, 118, 119, 127, 132, 430, 438; flagellaris × frondosus, 118, 119; flagellaris × hispidus, 126; flagellaris × procumbens, 129; floricomus, 106; floridus, 438; frondosus, 106, 107, 119, 122, 123, 427–429, 433–435; frondosus × hispidus, 120; frondosus montanus, 428; frondosus × nigricans, 119; frondosus × procumbens, 121; fruticosus, 427, 428, 430; geophilus, 128, 130; grandicaulis, 437; hispidus, 108, 124, 125, 127, 426–428, 430, 432–438; hispidus major, 436; hispidus × nigricans, 124; hispidus × procumbens, 111, 124; hispidus suberectus, 436; Millspaughii, 426–429; multiformis, 437; multispinus, 121; nigricans, 107, 108, 124, 126, 436; nigrobaccus, 106, 427, 429; nitidus, 439; obovalis, 430; occidentalis, 105; parvifolius, 434; philadelphicus, 435; procumbens, 109, 111, 116, 121, 122, 128, 131, 426, 431, 433, 438; procumbens roribaccus, 127; pubescens, 432; Randii, 437; recurvans, 107, 426, 433; recurvicaulis, 437; roribaccus, 128, 131; semisetosus, 436; sempervirens, 430; setosus, 431, 435, 436; strigosus, 105, 238, 242; suberectus, 427, 429; triflorus, 105, 432; trivialis, 426, 428, 431–433, 439; vermontanus, 125, 436; villosus, 427–432, 435; villosus Randii, 433
Rusby, H. H. Two new species of edible fruits from Mexico, 145
Rydberg, P. A. Notes on Rosaceae, 79, 351; Studies on the Rocky Mountain flora, 11
Rydbergia grandiflora, 238
Rynchospora dodecandra, 516

Sabal glabra, 227, 229, 231, 235; Palmetto, 232, 235
Sabalites, 405; **magothiensis,** 405
Sabbatia decandra, 232
Sabdariffa, 92–94; Alia, 92; Lobellij, 92
Salicaceae, 336
Salix, 334, 420–422; brachycarpa, 238, 240, 242; flexuosa, 421; fluviatilis, 335, 336; laevigata, 335, 336; Lesquereuxii,

409, 421; nigra, 227; pseudo-Hayei, 421; saximontana, 238
Santa Clara formation, 329
Sapindus, 423; Morrisoni, 421
Sargassum-?, 496; Agardhianum, 496; Filipendula, 496; Liebmanni, 496
Sassafras, 413–415; acutilobum, 409, 414, 415; cretaceum, 409, 412–415, 417; cretaceum dentatum, 414, 416; cretaceum obtusum, 414–416; cretaceum recurvatum, 412; integrifolium, 414; mirabile, 412, 413, 415; Mudgei, 414; obtusum, 414; recurvatum, 411, 412; subintegrifolium, 414
Saxifraga austromontana, 238; cernua, 238; chrysantha, 238; debilis, 238; flagellaris, 238; rhomboidea, 238
Saxosae, 80
Scinaia, 500; furcellata, 502; furcellata complanata, 500, 502; furcellata undulata, 500, 502; **latifrons,** 500, 502, 512
Scirpus Eriophorum, 229
Scoliosorus, 171
Scorzonera, 16
SCRIBNER, F. LAMSON-. Notes on certain species of Muhlenbergia, 319
Sedum, 313; stenopetalum, 238
Selaginella apus, 76; densa, 238, 240; rupestris, 78
Sematophyllum pungens, 36
Senecio atratus, 238; carthamoides, 238; chloranthus, 238, 240; crocatus, 238; eremophilus, 238, 240; Fendleri, 238; pudicus, 238; taraxicoides, 238; werneriaefolius, 238, 240
Sequoia, 341; Reichenbachii, 421, 422; sempervirens, 335, 336, 342
Serenoa serrulata, 231, 235, 516
Seseli, 14
SHERFF, E. E. A new variety of Carex lupulina, 481
Sibbaldia, 357; procumbens, 239
Sibbaldiopsis, 80, 362
Sieversia turbinata, 239
Significance of Monogramma, 185
Silene acaulis, 239
Sinapis, 478
Sisyrinchium grandiflorum, 376
Smilax auriculata, 232; lanceolata, 229; laurifolia, 225, 231; pumila, 516
Solidago decumbens, 239
Solorina crocea, 291; saccata spongiosa, 291
Some common species of Crataegus at Thompson's Mills, Georgia, 25
Sophia incisa, 239, 240
SOTH, B. List of plants collected above timber line on Pikes Peak, with altitudinal extensions and notes, 237
Sphaerium sp., 331
Sphaerophoraceae, 287
Sphaerophorus coralloides, 287; fragilis, 287
Spiraea latifolia, 447; tomentosa, 447
Spirogyra communis, 299; crassa, 299
Spirogyra, Conjugation of two different species of, 299
Stellaria crassifolia, 239, 240; umbellata, 239
Stephanomeria, 11; tenuifolia, 12; pauciflora, 12
Stereocaulon, 287; paschale, 289; tomentosum, 289; tomentosum simplex, 289
Stereophyllum **Howei,** 35
Sterigmatocystis niger, 137
STEVENS, N. E. Dioecism in the trailing arbutus, with notes on the morphology of the seed, 531
Sticta pulmonaria linita, 291
Stictolejeunea, 270
Stramonium, 94
Strawberry, pine, 361
Strobilites, 420
Studies on the Rocky Mountain flora—XXV, 11
Stylopappus grandiflorus, 18; laciniatus, 18, 20, 21; laciniatus longifolius, 21
Subjugae, 83
Suffruticulosae, 356
Supplemental description of Euphorbia ephedromorpha, A, 343
Swertia palustris, 239
Sympetalae, 540
Synthyris alpina, 239
Syrrhopodon Bernoullii, 33; brachysteloides, 33

Taravalia, 295–297; aptera, 296
Taraxacum, 21; angustifolium, 21; dumetorum, 21, 239, 240; leiospermum, 21
Taxilejeunea **obtusangula,** 215, 217, 218, 222
Taxithelium planum, 36
Taxodium, 235; distichum, 227–231, 235; imbricarium, 225, 226, 228, 231, 235
Telesonix Jamesii, 237
Teratological fruits of Ptelea, 385
Tetraneuris, 16; lanata, 239
Thalictrum alpinum, 239, 240; purpurascens, 382
Thelotrema lepadinum, 290
Thinnfeldia, 403, 404; rhomboidalis, 404
Thlaspi coloradense, 239
Thyrsopteris, 409
Tillandsia tenuifolia, 516; usneoides, 225–227, 229, 231, 234, 516, 518
Tonestus pygmaeus, 239
Tragopogon Dandelion, 16; lanatum, 16; virginicum, 16
Trautvetteria carolinensis, 232
Trichostelium fluviale, 36
Trifolium dasyphyllum, 239; nanum, 239

Trilisa odoratissima, 226
Trionum, 95
Trisetum subspicatum, 239
Troximon, 16, 18; arachnoideum, 18, 21; arizonicum, 19; aurantiacum, 19, 237; elatum, 19; glaucum, 17; glaucum dasycephalum, 17; glaucum laciniatum, 18, 21; gracilens, 18, 19; gracilens Greenei, 19; grandiflorum, 18, 20, 21; grandiflorum laciniatum, 21; grandiflorum obtusifolium, 20, 21; grandiflorum tenuifolium, 20, 21; laciniatum, 17, 18, 20; montanum, 20; pumilum, 17; purpureum, 18, 19; roseum, 17; villosum, 17
Trypethelium cruentum, 516
Two new species of edible fruits from Mexico, 145
Two new species of Opuntia, 141

Ulmus sp., 227
Ulvâ decorticata, 494; fasciata, 490; Lactuca, 490; Lactuca rigida, 490; Linza, 490; rigida, 490
Umbilicaria hyperborea, 291; Muhlenbergii alpina, 291; pustulata, 291
Umbilicariaceae, 291
Upon assimilation of atmospheric nitrogen by fungi, 135
Urceolaria scruposa, 290
Uredineae, New species of, 369
Uromyces **aemulus,** 373, 375; **agnatus,** 378; ambiguus, 374, 376; aterrimus, 374; aureus, 374, 375; bicolor, 374, 376; Hordei, 377; Houstoniatus, 376; Jatrophae, 378; **major,** 377; minimus, 377; **mysticus,** 377; oaxacanus, 378; primaverilis, 376; **probus,** 376; reticulatus, 376; Sisyrinchii, 376
Usnea plicata, 292
Usneaceae, 291

Vaccinium vacillans, 433
Vaccinioideae, 540
Valvata virens, 331
Vaseya comata, 319
Veronica alpina, 239, 240
Viburnum, 339
Viola affinis, 1, 2; amorphophylla, 3, 4; arvensis, 197, 198; asarifolia, 195; blanda, 7–9; Brainerdii, 8; Brittoniana, 5, 6, 9; canadensis, 1, 191, 194, 197; chalcosperma, 2; clandestina, 195; conspersa, 195–197; cucullata, 2, 3, 7; cucullata × fimbriatula, 3, 4; debilis, 196; dentata, 4, 5; denticulosa, 7; Egglestonii, 9; emarginata, 4, 5, 7; eriocarpa, 194, 197; esculenta, 2, 9; fimbriatula, 2–5, 7, 194; fimbriatula × palmata, 3; hastata, 191–193, 197; hastata glaberrima, 192; hirsutula, 192; incognita, 7–9; incognita **Forbesii,** 8; insignis, 6; labradorica, 195; lanceolata, 7; Langloisii, 1; Langloisii **pedatiloba,** 2; lobata, 192; multicaulis, 196; odorata, 196; pallens, 7; palmata, 3, 7; pedata, 6, 9; pedatifida, 6, 7, 9; Porteriana, 4, 5; primulifolia, 5, 7; primulifolia australis, 7; primulifolia villosa, 7; pubescens, 1, 194, 197; punctata, 195, 196; Rafinesquii, 197; renifolia, 7, 8; repens, 195; rostrata, 196, 197; rotundifolia, 195; sagittata, 2, 4, 5, 7, 194; sagittata emarginata, 5; scabriuscula, 1, 194; septemloba, 5–7, 9; sororia, 195; striata, 192–194, 196, 197; tenuipes, 193; tripartita, 191, 192, 197; tripartita glaberrima, 192–194, 197; uliginosa, 195; viarum, 2, 7; villosa, 2, 7, 192; vittata, 7; Walteri, 192, 196, 197
Vitis rotundifolia, 516
Vittaria, 154–190; elongata, 189; Gardneriana, 166; intramarginalis, 165, 170, 176, 178, 180, 187, 189, 190; lineata, 165, 166, 177, 178; **minima,** 164–166, 178, 188; minor, 165; Orbignyana, 166; pusilla, 165; remota, 163, 166, 176–178, 180, 183, 187–190, 275; scolopendrina, 189; scabrida, 188; sikkimensis, 164, 178–180, 189; stipitata, 166, 188
Vittarieae, 153, 158; genera, 158; general considerations, 182; ontogenetic stages of, 176; pluriseriate genera of, 167; tribal characters, 155
Vittarieae: their external morphology, venation, and relationships, The genera of the fern tribe, 153

WESTER, P. J. Contributions to the history and bibliography of the roselle, 91
Widdringtonites subtilis, 421, 422
WILLIAMS, R. S. Panama mossses, 33
WILSON, P. Notes on Rutaceae—V. Species characters in Ptelea and Taravalia, 295
WOODBURN, W. L. Development of the embryo sac and endosperm in some seedless persimmons, 379
Woodsia ilvensis, 72; obtusa, 73

Yukon region, List of lichens collected in the, 287

Zizania aquatica, 227, 229
Zizia cordata, 232
Zygadenus elegans, 239

EIGHT CUT-LEAVED SPECIES OF VIOLA, 1–4, palmate; 5–8, pedate.

U of M



BENEDICT: VITTARIEAE, TRIBAL CHARACTERS

UofM



BENEDICT: VITTARIEAE, MONOGRAMMA



BENEDICT: VITTARIEAE, HECISTOPTERIS



BENEDICT: VITTARIEAE, VITTARIA



U of M

BENEDICT: VITTARIEAE, POLYTAENIUM and ANTROPHYUM

BENEDICT: VITTARIEAE, POLYTAENIUM and ANTROPHYUM



BENEDICT: VITTARIEAE, ONTOGENETIC STAGES



1—12 RECTOLEJEUNEA BRITTONIAE Evans

13—19 CERATOLEJEUNEA INTEGRIFOLIA Evans

UofM

HELIOTYPE CO. BOSTON



TAXILEJEUNEA OBTUSANGULA (Spruce) Evans



1—8 LEPTOCOLEA SCABRIFLORA (Gottsche) Evans

9—16 LEPTOCOLEA PLANIFOLIA Evans

HELIOTYPE CO. BOSTON



1–3 LEPTOCOLEA CARDIOCARPA (Mont.) Evans

4—10 APHANOLEJEUNEA EXIGUA Evans

11—16 APHANOLEJEUNEA CRENATA Evans

18—26 APHANOLEJEUNEA SICAEFOLIA (Gottsche) Evans

HELIOTYPE CO. BOSTON



M.A.K. del.ad nat.

KINGSLEY, RHIZOME AND ROOT OF DELPHINIUM SCAPOSUM



Fig. 7.

M.A.K del. ad. nat.

KINGSLEY, RHIZOME OF DELPHINIUM SCAPOSUM

1



2

3



4

5



6

7

8

9

10

HANNIBAL, PLIOCENE FLORA

M 701



WOODBURN, EMBRYO SAC AND ENDOSPERM OF SEEDLESS PERSIMMONS



HARRIS, TERATOLOGICAL FRUITS OF PTELEA

U of M



UPPER CRETACEOUS FILICALES

UofM



UPPER CRETACEOUS DICOTYLEDONES



GAGER: ONAGRA BIENNIS, 15A BROAD

GAGER: ONAGRA BIENNIS, 15A NARROW

MacDOUGAL, OPUNTIA ON SAHUARO



MACDOUGAL, OPUNTIA ON SAHUARO



MACDOUGAL, OPUNTIA ON ACACIA

MacDOUGAL, OPUNTIA ON ACACIA



CAREX LUPULINA ALBOMARGINATA Sherff



DICTYOTA VIVESII M. A. HOWE

SCINAIA LATIFRONS M. A. HOWE



ANATHECA DICHOTOMA M. A. HOWE

GRACILARIA VIVESII M. A. HOWE

Mrau



FAUCHEA SEFFERI M. A. HOWE

FAUCHEA (?) MOLLIS M. A. HOWE



1



2



3

4



5

6



7

1–5. GRACILARIA VIVESII M. A. HOWE
6. FAUCHEA (?) MOLLIS M. A. HOWE
7. CLADOPHORA MACDOUGALII M. A. HOWE



HALYMENIA ACTINOPHYSA M. A. HOWE

CYATHEA MULTIFLORA J. E. SMITH

VOL. 38 DECEMBER, 1911 NO. 12

GENERAL LIBRARY
UNIVERSITY OF
JAN 17

BULLETIN

OF THE

TORREY BOTANICAL CLUB

Editor

PHILIP DOWELL

Associate Editors

JOHN HENDLEY BARNHART
JEAN BROADHURST
ERNEST DUNBAR CLARK
ALEXANDER WILLIAM EVANS
TRACY ELLIOT HAZEN
MARSHALL AVERY HOWE
HERBERT MAULE RICHARDS
NORMAN TAYLOR

CONTENTS

Dioecism in the trailing arbutus, with notes on the morphology of the seed . . . NEIL E. STEVENS 531

On the identity of Cyathea multiflora, type of the genus Hemitelia R. Br. (Plate 35) . . . . . WILLIAM R. MAXON 545

INDEX TO AMERICAN BOTANICAL LITERATURE 551

INDEX TO VOLUME 38 559

PUBLISHED FOR THE CLUB

THE NEW ERA PRINTING COMPANY

LANCASTER, PA.

THE TORREY BOTANICAL CLUB

President,
HENRY H. RUSBY, M.D.

Vice-Presidents,
EDWARD S. BURGESS, PH.D. JOHN HENDLEY BARNHART, A.M., M.D

Secretary and Treasurer,
BERNARD O. DODGE, PH. B.,
Dept. of Botany, Columbia University,
New York City.

MEETINGS

Meetings twice each month from October to May inclusive: the second Tuesday, at 8:00 P.M., at the American Museum of Natural History; the last Wednesday, at 3:30 P.M., in the Museum Building of the New York Botanical Garden.

PUBLICATIONS

All subscriptions and other business communications relating to the publications of the Club should be addressed to the Treasurer, Bernard O. Dodge, Dept. of Botany, Columbia University, New York City.

Bulletin. Monthly, established 1870. Price, $3.00 a year; single numbers 30 cents. Of former volumes, only 24–37 can be supplied separately; certain numbers of other volumes are available, but the entire stock of some numbers has been reserved for the completion of sets. Manuscripts intended for publication in the BULLETIN should be addressed to Philip Dowell, Editor, Port Richmond, N. Y.

Torreya. Monthly, established 1901. Price, $1.00 a year. Manuscripts intended for publication in TORREYA should be addressed to Norman Taylor, Editor, Central Museum, Eastern Parkway, Brooklyn, N. Y.

Memoirs. Occasional, established 1889. (See last pages of cover.)

Preliminary Catalogue of Anthophyta and Pteridophyta within 100 miles of New York City, 1888. Price, $1.00

Reprints. Contributors of articles accepted for publication in the BULLETIN should order reprints, if desired, when they return galley proof to the editor. 25 copies of reprints without cover may be had gratis. If cover is wanted, or if more than 25 copies are wanted, they may be ordered at the following rates from The New Era Printing Co., Lancaster, Pa., the contributor paying for all.

	2pp	4pp	8pp	12pp	16pp	20pp
25 copies	$ .75	$1.05	$1.30	$1.80	$2.20	$2.50
50 copies	.90	1.20	1.70	2.20	2.50	2.85
100 copies	1.15	1.55	1.95	2.55	2.90	3.20
200 copies	1.70	2.35	2.90	3.75	4.35	4.70

Covers: 25 for 75 cents, additional covers 1 cent each.

Plates for reprints, 40 cents each per 100.

American Fern Journal

An illustrated quarterly devoted to the general study of ferns. Subscription, including membership in the AMERICAN FERN SOCIETY, $1.00 a year, should be sent to H. G. RUGG, Treasurer, HANOVER, N. H. Matter for publication should be addressed to PHILIP DOWELL, PORT RICHMOND, N. Y.

THE BRYOLOGIST

WITH THE JANUARY, 1911, NUMBER BEGINS ITS FOURTEENTH YEAR AND VOLUME

It is a 16–20 page bi-monthly devoted to the study of the Mosses, Hepatics, and Lichens. It is fully illustrated with new, original, and artistic drawings and half-tones. It is indispensable to the working bryologist, professional as well as amateur. Send for sample copy. Subscription price $1.25 a year.

Address, **Dr. A. J. GROUT, New Dorp, N. Y.**

INDEX TO AMERICAN BOTANICAL LITERATURE

This Index, printed each month in the BULLETIN, is reissued on library catalog cards; these are furnished to subscribers at the rate of one cent for each card.

Subscriptions to the card issue may be addressed to the Treasurer of the Torrey Botanical Club,

BERNARD O. DODGE, DEPT. OF BOTANY, Columbia University, N. Y. City.

From January, 1903, to December, 1909, inclusive, the Index was also reprinted on paper, on one side only, and can be furnished in that form at the rate of $3.00 a year.

Back Numbers of The

BULLETIN OF THE TORREY BOTANICAL CLUB

are wanted to complete files as follows:

Vol. 7 (1880), Nos. 1 and 9
Vol. 9 (1882), No. 1
Vol. 11 (1884), Nos. 1 and 2
Vol. 12 (1885), Nos. 2 and 3
Vol. 15 (1888), No. 2
Vol. 16 (1889), No. 4
Vol. 17 (1890), No. 4
Vol. 18 (1891), No. 1

Also, MEMOIRS, Vol. 9.

Those possessing the numbers desired are requested to state prices. Address

TREASURER, TORREY BOTANICAL CLUB

Dept. of Botany, Columbia University, New York City

North American Flora

THIS work is designed to present descriptions of all plants growing independently of cultivation, in North America, here taken to include Greenland, Central America, the Republic of Panama, and the West Indies, except Trinidad, Tobago, and Curaçao and other islands off the north coast of Venezuela.

It will be published in parts at irregular intervals by the New York Botanical Garden through the aid of the income of the David Lydig Fund bequeathed by Charles P. Daly.

It is planned to issue parts as rapidly as they can be prepared, the extent of the work making it possible to commence publication at any number of points. The completed work will form a series of volumes with the following sequence :

Volume 1. Mycetozoa, Schizophyta, Diatomaceae.
Volumes 2–10. Fungi.
Volumes 11–13. Algae.
Volumes 14 and 15. Bryophyta.
Volume 16. Pteridophyta, Gymnospermae.
Volumes 17–19. Monocotyledones.
Volumes 20–32. Dicotyledones.

The preparation of the work has been referred by the Scientific Directors of the Garden to a committee consisting of Dr. N. L. Britton, Dr. W. A. Murrill, and Dr. J. H. Barnhart.

Professor George F. Atkinson, of Cornell University, Professors Charles R. Barnes and John M. Coulter, of the University of Chicago, Mr. Frederick V. Coville, of the United States Department of Agriculture, Professor Edward L. Greene, of the United States National Museum, Professor Byron D. Halsted, of Rutgers College, and Professor William Trelease, of the Missouri Botanical Garden, have consented to act as an advisory committee.

The following parts have been published :

Vol. 3. Part 1, Nectriaceae, Hypocreaceae, by F. J. Seaver ; Chaetomiaceae, by H. L. Palliser ; Fimetariaceae, by D. Griffiths and F. J. Seaver.

Vol. 7. Part 1, Ustilaginales, by G. P. Clinton. Part 2, Coleosporiaceae, Uredinaceae, Aecidiaceae (pars), by J. C. Arthur.

Vol. 9. Parts 1 and 2, Polyporaceae, by W. A. Murrill. Part 3, Boletaceae, by W. A. Murrill ; Agaricaceae (pars), by W. A. Murrill and G. S. Burlingham.

Vol. 16. Part 1, Ophioglossaceae, by L. M. Underwood and R. C. Benedict ; Marattiaceae, by L. M. Underwood; Osmundaceae, Ceratopteridaceae, by R. C. Benedict ; Schizaeaceae, Gleicheniaceae, Cyatheaceae (pars), by W. R. Maxon.

Vol. 17. Part 1, Typhaceae, by P. Wilson; Sparganiaceae, Elodeaceae, Hydrocharitaceae, by P. A. Rydberg ; Zannichelliaceae, Zosteraceae, Cymodoceaceae, Naiadaceae, Lilaeaceae, by N. Taylor ; Scheuchzeriaceae, by N. L. Britton ; Alismaceae, by J. K. Small ; Butomaceae, Poaceae (pars), by G. V. Nash.

Vol. 22. Part 1, Podostemonaceae, by G. V. Nash ; Crassulaceae, by N. L. Britton and J. N. Rose ; Penthoraceae, Parnassiaceae, by P. A. Rydberg. Part 2, Saxifragaceae, Hydrangeaceae, by J. K. Small and P. A. Rydberg; Cunoniaceae, Iteaceae, Hamamelidaceae, by N. L. Britton ; Pterostemonaceae, by J. K. Small ; Altingiaceae, by P. Wilson ; Phyllonomaceae, by H. H. Rusby. Part 3, Grossulariaceae, by F. V. Coville and N. L. Britton ; Platanaceae, by H. A. Gleason ; Crossosomataceae, by J. K. Small ; Connaraceae, by N. L. Britton ; Calycanthaceae, by C. L. Pollard ; Rosaceae (pars), by P. A. Rydberg. Part 4, Rosaceae (pars), by P. A. Rydberg.

Vol. 25. Part 1, Geraniaceae, by L. T. Hanks and J. K. Small ; Oxalidaceae, Linaceae, by J. K. Small ; Erythroxylaceae, by N. L. Britton. Part 2, Tropaeolaceae, by G. V. Nash ; Balsaminaceae, Limnanthaceae, by P. A. Rydberg ; Koeberliniaceae, by J. H. Barnhart ; Zygophyllaceae, by A. M. Vail and P. A. Rydberg ; Malpighiaceae, by J. K. Small.

The subscription price is fixed at $1.50 for each part ; it is expected that four or five parts will be required for each volume. A limited number of separate parts (excepting Part 1, Vol. 9) will be sold at $2.00 each. Address

THE NEW YORK BOTANICAL GARDEN
BRONX PARK, NEW YORK CITY

PUBLICATIONS BY OFFICERS AND STUDENTS OF THE DEPARTMENT OF BOTANY, COLUMBIA UNIVERSITY

1. MEMOIRS OF THE DEPARTMENT:

Vol. 1. A Monograph of the North American Species of the Genus *Polygonum* (1895). By John Kunkel Small, Fellow in Botany, 1893–1895; Curator of the Herbarium, 1895–1898.

Quarto, 178 pages, 84 plates. Price $6.00.

Vol. 2. A Monograph of the North American Potentilleae (1898). By Per Axel Rydberg, Fellow in Botany, 1896–7.

Quarto, 224 pages, 112 plates. Price $6.00.

2. CONTRIBUTIONS FROM THE DEPARTMENT:

Vol. 1. Nos. 1–25. 1886–1892. Price $5.00.
Vol. 2. Nos. 26–50. 1892–1894. Price $5.00.
Vol. 3. Nos. 51–75. 1894–1895. Price $5.00.
Vol. 4. Nos. 76–100. 1895–1896. Price $5.00.
Vol. 5. Nos. 101–125. 1896–1897. Price $5.00.
Vol. 6. Nos. 126–150. 1897–1898. Price $5.00.
Vol. 7. Nos. 151–175. 1898–1901. Price $5.00.
Vol. 8. Nos. 176–200. 1901–1902. Price $5.00.
Vol. 9. Nos. 201–. 1903–(current).

List of separate numbers available on application.

3. A Text-book of General Lichenology (1896). By Albert Schneider, Fellow in Botany, 1895–1896.

Octavo, 230 pages, 76 plates. Price $4.25 (cloth); $3.80 (paper). Published by Willard N. Clute & Co. Binghamton, N. Y.

The Department invites propositions relative to exchanges of herbarium material for any of the above named publications. Address Dr. Carlton C. Curtis, Columbia University, N. Y. City.

4. An Illustrated Flora of the Northern United States, Canada, and the British Possessions from Newfoundland to the Parallel of the Southern Boundary of Virginia, and from the Atlantic Ocean westward to the 102d Meridian (1896–1898). By Nathaniel Lord Britton, Professor of Botany, 1891–1896; Emeritus Professor 1896–, and Hon. Addison Brown.

Three volumes, royal octavo. Vol. 1, 612 pages; Vol. 2, 642 pages; Vol. 3, 588 pages; 4162 figures in the text, illustrating every species described.

Published by Charles Scribner's Sons, New York. Price, in cloth, $9.00 for the three volumes; with the indexes and keys bound separately, $10.00.

5. Our Native Ferns and their Allies. (Sixth Edition) 1900. By Lucien Marcus Underwood, Professor of Botany, 1896–1907.

Duodecimo, 158 pages, 35 figures. Published by Henry Holt & Co., New York. Price, $1.00.

6. Moulds, Mildews, and Mushrooms. By Lucien Marcus Underwood, Professor of Botany, 1896–1907.

Duodecimo, 236 pages, 10 plates. Published by Henry Holt & Co. Price $1.50.

7. Nature and Development of Plants (1907). By Carlton Clarence Curtis, Assistant in Botany, 1892–1895; Tutor in Botany, 1895–1906; Instructor in Botany, 1906–1908; Adjunct Professor of Botany, 1908–1910; Associate Professor of Botany, 1910–.

Octavo, 471 pages, 42 illustrations. Published by Henry Holt & Co., New York. Price $2.50.

ESTABLISHED 1851

EIMER & AMEND

205–211 Third Ave., Corner 18th Street NEW YORK

MANUFACTURERS AND IMPORTERS OF

Chemicals and Chemical Apparatus

Our Specialties

MICROSCOPES and ACCESSORIES

MICROTOMES

BALANCES and WEIGHTS

Sole Agents for

C. A. F. KAHLBAUM'S C. P. Chemicals and Reagents

CARL SCHLEICHER & SCHÜLL'S C. P. Filters

SCHOTT & GENOSSEN'S Famous Laboratory Glassware

And for many other renowned European houses in our line.

COMPLETE LINE OF

Bacteriological Apparatus

WE HANDLE

EVERYTHING NEEDED FOR A LABORATORY

Inspection of premises cordially invited.

Vol. 5; price, $3.00:

List of *Pteridophyta* and *Spermatophyta* growing without cultivation in northeastern North America. Pages 1–377. 4 D 1893–31 D 1894.

Vol. 6, No. 1; price, $1.25:

Rusby, Henry Hurd. An enumeration of the plants collected in Bolivia by Miguel Bang. Part 3. Pages 1–130. 17 N 1896.

Vol. 6, No. 2; not furnished separately:

Grout, Abel Joel. A revision of the North American *Sotheciaceae* and *Brachythecia*. Pages 131–210. 30 Jl 1897.

Vol. 6, No. 3; price, 50 cents:

Hazen, Tracy Elliot. The life history of *Sphaerella lacustris* (*Haematococcus pluvialis*). Pages 211–246, *plates 86, 87* (*colored*). 8 Je 1899.

Vol. 6, No. 4; price, 50 cents:

Underwood, Lucien Marcus. A review of the genera of ferns proposed prior to 1832. Pages 247–283. 1 D 1899.

Vol. 6, No. 5; price, 25 cents:

Fink, Bruce. Notes on lichen distribution in the upper Mississippi Valley. Pages 285–307. 1 D 1899.

Vol. 7; price, $3.00:

Howe, Marshall Avery. The *Hepaticae* and *Anthocerotes* of California. Pages 1–208, *plates 88–122*. 5 Au 1899.

Vol. 8, No. 1; not furnished separately:

Lloyd, Francis Ernest. The comparative embryology of the *Rubiaceae*. Part I. Pages 1–26, *plates 1–4*. 26 Au 1899. Part II. Pages 27–112, *plates 5–15* 15 F 1902.

Vol. 8, No. 2; price, $1.00:

Evans, Alexander William. The *Lejeuneae* of the United States and Canada. Pages 113–183, *plates 16–22*. 15 F 1902.

Vol. 8, No. 3; price, 75 cents:

Britton, Elizabeth Gertrude, & Taylor, Alexandrina. The life history of *Vittaria lineata*. Pages 185–211, *plates 23–31*. 30 Au 1902.

Vol. 9; price, $3.00:

Salmon, Ernest Stanley. A monograph of the *Erysiphaceae*. Pages 1–292, *plates 1–9*. 4 O 1900. No longer sold except in complete sets of the Memoirs.

Vol. 10; price, $3.00:

Burgess, Edward Sandford. History of Pre-Clusian botany in its relation to *Aster*. Pages i–xii, 1–447. 22 N 1902.

Vol. 11, No. 1; price, $1.75:

Griffiths, David. The North American *Sordariaceae*. Pages 1–134, *plates 1–19*. 30 My 1901.

Vol. 11, No. 2; price, $1.75:

Hazen, Tracy Elliot. The *Ulothricaceae* and *Chaetophoraceae* of the United States. Pages 135–250, *plates 20–42*. 20 O 1902.

Vol. 12, No. 1; price, $1.50:

Northrop, Alice Rich. Flora of New Providence and Andros (Bahama Islands). Pages 1–98, *plates 1–19 + map*. 10 D 1902.

Vol. 12, No. 2; price, $1.00:

Banker, Howard James. A contribution to a revision of the North American *Hypnaceae*. Pages 99–194. 13 Je 1906.

Vol. 12, No. 3; price, 75 cents:

Kupfer, Elsie. Studies in plant regeneration. Pages 195–241. 10 Je 1907.

Vol. 13; price, $3.00:

Burgess, Edward Sandford. Species and variations of Biotian Asters, with discussion of variability in *Aster*. Pages i–xv, 1–419, *plates 1–13*. 15 Mr 1906.

Vol. 14, No. 1; price, $1.25:

Burlingham, Gertrude Simmons. A study of the *Lactariae* of the United States. Pages 1–109, *figures 1–15*. 26 My 1908.

Vol. 14, No. 2; price, 75 cents:

Butler, Ormond. Observations on the California vine disease. Pages 111–153, *plates 1–5*. 18 Je 1910.

Address: **BERNARD O. DODGE,**
Dept. of Botany. Columbia University. New York City.

MEMOIRS OF THE TORREY BOTANICAL CLUB

A series of technical papers on botanical subjects, published at irregular intervals. Price $3.00 a volume. Not offered in exchange.

Vol. 1, No. 1; not furnished separately:

Bailey, Liberty Hyde. Studies of the types of various species of the genus *Carex*. Pages 1–85. 25 My 1889.

Vol. 1, No. 2; not furnished separately:

Martindale, Isaac Comly. Marine algae of the New Jersey coast and adjacent waters of Staten Island. Pages 87–111. 24 Au 1889.

Vol. 1, No. 3; price, 75 cents:

Spruce, Richard. *Hepaticae* Bolivianae, in Andibus Boliviae orientalis, annis 1885–6, a cl. H. H. Rusby lectae. Pages 113–140. 20 Ja 1890.

Vol. 1, No. 4; price, 75 cents:

Sturtevant, Edward Lewis. On seedless fruits. Pages 141–187. 30 My 1890.

Vol. 2, No. 1; not furnished separately:

Halsted, Byron David. Reserve food-materials in buds and surrounding parts. Pages 1–26, *plates 1, 2*. 10 S 1890.

Vol. 2, No. 2; price, 75 cents:

Vail, Anna Murray, & Hollick, Charles Arthur. Contributions to the botany of Virginia. Pages 27–56, *plates 3, 4*. 23 D 1890.

Vol. 2, No. 3; not furnished separately:

Holm, Herman Theodor. Contributions to the knowledge of the germination of some North American plants. Pages 57–108, *plates 5–19*. 15 Ap 1891.

Vol. 2, No. 4; price, 75 cents:

Wheelock, William Ebner. The genus *Polygala* in North America. Pages 109–152. 30 D 1891.

Vol. 3, No 1; not furnished separately:

Small, John Kunkel, & Heller, Amos Arthur. Flora of western North Carolina and contiguous territory. Pages 1–39. 20 F 1892.

Vol 3, No. 2; price, $2.00:

Morong, Thomas. The *Naiadaceae* of North America [with illustrations of all the species]. Pages 1–65, *plates 20–74*. 15 Mr 1893.

Vol. 3, No. 3; not furnished separately:

Rusby, Henry Hurd. An enumeration of the plants collected in Bolivia by Miguel Bang. [Part 1.] Pages 1–67. 28 Ap 1893.

Vol. 4, No. 1; not furnished separately:

Underwood, Lucien Marcus. Index *Hepaticarum*. Part I, Bibliography. Pages 1–91. 10 Je 1893.

Vol. 4, No. 2; not furnished separately:

Small, John Kunkel, & Vail, Anna Murray. Report on the botanical exploration of southwestern Virginia during the season of 1892. Pages 93–201, *plates 75–82*. 18 N 1893–17 Ap 1894.

Vol. 4, No. 3; price, 50 cents:

Rusby, Henry Hurd. An enumeration of the plants collected in Bolivia by Miguel Bang. Part 2. Pages 203–274. 10 Mr–17 Ap 1895.

Vol. 4, No. 4; price, 50 cents:

Pettit, Anna Stockton. *Arachis hypogaea* L. Pages 275–296, *plates 83–85*. 5 Je 1895.

Vol. 4, No. 5; price, 75 cents:

Rydberg, Per Axel. The North American species of *Physalis* and related genera. Pages 297–374. 15 S 1896.